

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
29 March 2001 (29.03.2001)

PCT

(10) International Publication Number
WO 01/21577 A2(51) International Patent Classification⁷: C07C 235/00

(21) International Application Number: PCT/JP00/06375

(22) International Filing Date:
19 September 2000 (19.09.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
11/266298 20 September 1999 (20.09.1999) JP
11/357889 16 December 1999 (16.12.1999) JP
2000/126272 20 April 2000 (20.04.2000) JP(71) Applicant (for all designated States except US): TAKEDA
CHEMICAL INDUSTRIES, LTD. [JP/JP]; 1-1,
Doshomashi 4-chome, Chuo-ku, Osaka-shi, Osaka
541-0045 (JP).


(72) Inventors; and

(75) Inventors/Applicants (for US only): KATO, Kaneyoshi
[JP/JP]; 2-40, Maruyamada 2-chome, Kawanishi-shi,
Hyogo 666-0152 (JP). TERAUCHI, Jun [JP/JP]; 3-5-204,
Hachizuka 3-chome, Ikeda-shi, Osaka 563-0024 (JP).
MORI, Masaaki [JP/JP]; 7-9-702, Kasuga 1-chome,
Tsukuba-shi, Ibaraki 305-0821 (JP). SUZUKI, Nobuhiro
[JP/JP]; 1077-50, Oaza-yatabe, Tsukuba-shi, Ibaraki
305-0861 (JP). SHIMOMURA, Yukio [JP/JP]; 12-1-410,Matsushiro 3-chome, Tsukuba-shi, Ibaraki 305-0035
(JP). TAKEKAWA, Shiro [JP/JP]; 5-3-B305, Umezono
2-chome, Tsukuba-shi, Ibaraki 305-0045 (JP). ISHI-
HARA, Yuji [JP/JP]; 12-30-305, Ninomiya 1-chome,
Tsukuba-shi, Ibaraki 305-0051 (JP).(74) Agents: TAKAHASHI, Shuichi et al.; Osaka Plant of
Takeda Chemical Industries, Ltd., 17-85, Jusohonmachi
2-chome, Yodogawa-ku, Osaka-shi, Osaka 532-0024 (JP).(81) Designated States (national): AE, AG, AL, AM, AU, AZ,
BA, BB, BG, BR, BY, BZ, CA, CN, CR, CU, CZ, DM, DZ,
EE, GD, GE, HR, HU, ID, IL, IN, IS, JP, KG, KR, KZ, LC,
LK, LR, LT, LV, MA, MD, MG, MK, MN, MX, MZ, NO,
NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TR, TT, UA, US,
UZ, VN, YU, ZA.(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— Without international search report and to be republished
upon receipt of that report.For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: MELANIN CONCENTRATING HORMONE ANTAGONIST



further substituents; R¹ and R² are independently hydrogen atom or a hydrocarbon group which may have substituents; R¹ and R², together with the adjacent nitrogen atom, may form a nitrogen-containing hetero ring which may have substituents; R² may form a spiro ring together with Ar; or R², together with the adjacent nitrogen atom and Y, may form a nitrogen-containing hetero ring which may have substituents; or a salt thereof; which is useful as an agent for preventing or treating obesity, etc.

WO 01/21577 A2

WO 01/21577

PCT/JP00/06375

1

DESCRIPTION

Melanin Concentrating Hormone Antagonist

5 TECHNICAL FIELD

The present invention relates to a melanin-concentrating hormone antagonist which is useful as an agent for preventing or treating obesity, etc.

10 BACKGROUND ART

Feeding behavior is an essential action for many living beings including humans. Therefore, if irregularities in feeding behavior occur, disorders, often connected to diseases, will occur in normal life-maintaining activities. Accompanying recent changes of our dietary environment, obesity is now becoming a social problem. In addition, not only is obesity a serious risk factor for life-style diseases such as diabetes, hypertension, and arteriosclerosis; it is also widely known that increased body weight places excessive burdens on joints such as knee joints, causing arthritis and pain.

The "diet boom," etc. show that there is a potentially great percentage of the population hoping to reduce body weight; on the other hand, many cases of feeding problems such as overeating, occurring due to causes such as hereditary neurosis or neurosis due to stress, have been reported.

Therefore, research on and development of agents for preventing or treating obesity, or agents for inhibiting eating, have been vigorously done for a long time.

The centrally acting anorectic drug, Mazindol, is now being marketed.

Many appetite control factors such as leptin, have recently been discovered, and the development of anti-obesity agents or anorectic agents which will regulate the functions of these appetite control factors is progressing.

WO 01/21577


PCT/JP00/06375

2

In particular, it is known that melanin- concentrating hormone (hereinafter also abbreviated as "MCH") originates in the hypothalamus and has orexigenic action. In addition, it has been reported that even though the daily behavior of MCH knock-out mice was normal, the amount of feeding by MCH knock-out mice was significantly reduced and their body weights were lighter than those of normal mice [Nature, Vol. 396, p.670, 1998]. This indicates that, if a MCH antagonist was produced, it can be expected to be an excellent anorectic agent or anti-obesity agent; but at present there are no known compound, especially non-peptide type compounds, which possess MCH antagonistic actions.

On the other hand, the following compounds are known as amine derivatives.

1) W098/38156 describes a compound of the formula :



wherein Ar is an optionally substituted ring assembly aromatic group or an optionally substituted condensed aromatic group; X is a bond, etc.; Y is an optionally substituted bivalent C₁₋₆ aliphatic hydrocarbon group which may have an intervening oxygen atom or sulfur atom; R¹ and R² are independently hydrogen atom or a lower alkyl, or R¹ and R², together with the adjacent nitrogen atom, form an optionally substituted nitrogen-containing hetero ring; Ring A is a benzene ring which may have further substituents in addition to the groups of the formula : -X-Ar where each symbol has the same meaning as defined above; Ring B is a 4 to 8 membered ring which may have further substituents in addition to the group of the formula : -Y-NR¹R² where each symbol has the same meaning as defined above; with the proviso that the condensed ring formed by ring A and ring B is an indole ring, the group of the formula : -X-Ar where


WO 01/21577

PCT/JP00/06375

3


each symbol has the same meaning as defined above is substituted at the 4-, 6-, or 7- position on the indole ring; or its salt, which has an action of inhibiting the production and secretion of β -amyloid protein.

5 2) W095/32967 describes compound of the formula :



wherein A is CONR, in which R is hydrogen or C₁₋₆ alkyl; Q is an optionally substituted 5 to 7 membered hetero ring containing 1 to 3 hetero atoms selected from nitrogen or sulfur; R¹ is hydrogen, halogen, etc.; R² and R³ are independently hydrogen, halogen, etc.; R₄ and R₅ are independently hydrogen or C₁₋₆ alkyl; R⁶ is halogen, hydroxy, etc.; R₇ and R₈ are independently hydrogen, C₁₋₆ alkyls, etc.; m is 0 to 4; n is 0, 1 or 2; or its salt, which has 5HT1D antagonist activity and can be expected to ameliorate anorexia.

3) W098/15274 describes a compound of the formula :




wherein Ar is phenyl, etc.; X is -O- or -S-; Y is CR⁵R^{5'} where R^{5'} is H and R⁵ is -H, etc.; Z is -CH₂- or -N-; R is H or -(C1-C6) alkyl; R¹ and R² are independently -(C1-C6) alkyl, etc.; R³ is H etc.; R⁴ is hydrogen, etc.; m is an integer of 0 to 2; q is 0 or 1; n is an integer of 0 to 4; p is an integer of 1 to 6; t is an integer of 1 to 4; which has an anti-oxidant activity and can be expected to ameliorate Alzheimer's disease.

4) EP533266


WO 01/21577

PCT/JP00/06375

4



wherein R^1 is halogen, etc.; R^2 is phenyl optionally substituted by 1 or 2 substituents selected from halogen, etc.; R^3 is




5

; R^4 and R^5 are independently hydrogen, halogen, etc.; R^{11} is hydrogen or C_{1-6} alkyl; which has 5HT1D antagonist activity, and can be expected to ameliorate anorexia.

There has been great desire for the development of a melanin-concentrating hormone antagonist which is useful as an agent for preventing or treating obesity, excellent in oral absorbency, and safe.

DISCLOSURE OF INVENTION

As a result of intensive studies of compounds with a MCH antagonistic action, the present inventors found that a derivative which is obtained by introducing a group of the formula : Ar^1-X- where each symbol has the same meaning as defined hereafter, into a compound of the formula :



20

wherein each symbol has the same meaning as defined hereinafter, had an excellent MCH antagonistic actions, to complete this invention.

Namely, the present invention relates to :


(1) a melanin-concentrating hormone antagonist which comprises a compound of the formula :

25

WO 01/21577

PCT/JP00/06375

5



wherein Ar¹ is a cyclic group which may have substituents;
X is a spacer having a main chain of 1 to 6 atoms;
Y is a bond or a spacer having a main chain of 1 to 6 atoms;
Ar is a monocyclic aromatic ring which may be condensed with
a 4 to 8 membered non-aromatic ring, and may have further
substituents;

R¹ and R² are independently hydrogen atom or a hydrocarbon
group which may have substituents; R¹ and R², together with
the adjacent nitrogen atom, may form a nitrogen-containing
hetero ring which may have substituents; R² may form a spiro
ring together with Ar; or R², together with the adjacent
nitrogen atom and Y, may form a nitrogen-containing hetero
ring which may have substituents; or a salt thereof;

(2) an antagonist according to the above (1), wherein Y is
a spacer having a main chain of 1 to 6 atoms; R¹ and R² are
independently hydrogen atom or a hydrocarbon group which
may have substituents; R¹ and R², together with the adjacent
nitrogen atom, may form a nitrogen-containing hetero ring
which may have substituents; or R² may form a spiro ring
together with Ar;

(3) an antagonist according to the above (2), wherein Ar¹
is an aromatic group which may have substituents; and "a
hydrocarbon group which may have substituents" for R¹ and
R² is "C₁₋₆ alkyl which may have substituents";

(4) an antagonist according to the above (1), wherein the
cyclic group for Ar¹ is C₆₋₁₄ monocyclic or condensed
polycyclic aromatic hydrocarbon group;

(5) an antagonist according to the above (1), wherein the
cyclic group for Ar¹ is a group formed by removing an
optional one hydrogen atom from an aromatic ring assemble
in which 2 or 3 C₆₋₁₄ monocyclic or condensed polycyclic
aromatic hydrocarbon groups are directly bonded by single

WO 01/21577

PCT/JP00/06375

6

bonds;

(6) an antagonist according to the above (1), wherein the cyclic group for Ar¹ is a group formed by removing an optional one hydrogen atom from an aromatic ring assemble
5 in which C₆₋₁₄ monocyclic or condensed polycyclic aromatic hydrocarbon and 5 to 10 membered aromatic hetero ring are directly bonded by a single bond;

(7) an antagonist according to the above (1), wherein Ar¹ is phenyl, biphenyl, phenyl-pyridyl, phenyl-furyl,
10 phenyl-isoxazolyl, diphenyl-oxazolyl, pyridyl-phenyl, phenyl-pyrimidinyl, benzofuranyl-phenyl, furyl-phenyl, terphenyl, thienyl-phenyl, indolyl, naphthyl-oxadiazolyl, benzofuranyl-oxadiazolyl, benzothienyl, benzofuranyl, fluorenyl, pyridyl-pyrrolyl or
15 thioxanthanyl;


each of which may have 1 to 3 substituents selected from the group consisting of halogen atom; nitro; C₁₋₃ alkylenedioxy; optionally halogenated C₁₋₆ alkyl; hydroxy-C₁₋₆ alkyl; optionally halogenated C₃₋₆ cycloalkyl;
20 optionally halogenated C₁₋₆ alkoxy; optionally halogenated C₁₋₆ alkythio; hydroxy; C₇₋₁₉ aralkyloxy which may have substituents; C₆₋₁₄ aryloxy which may have substituents; amino; mono-C₁₋₆ alkylamino; di-C₁₋₆ alkylamino; 5 to 7 membered saturated cyclic amino which may have substituents
25 and may be condensed with a benzene ring; 5 to 7 membered non-aromatic heterocyclic groups which may have substituents; formyl; carboxy; C₆₋₁₄ aryl-carbonyl which may have substituents; C₆₋₁₄ aryl-carbamoyl which may have substituents; aromatic hetero ring-carbamoyl which may
30 have substituents; C₁₋₆ alkoxy-carbonyl; optionally halogenated C₁₋₆ alkyl-carboxamide; C₆₋₁₄ aryl-carboxamide which may have substituents; C₇₋₁₉ aralkyl-carboxamide which may have substituents; aromatic hetero ring-carboxamide which may have substituents; N-(C₆₋₁₄ aryl-carbonyl which
35 may have substituents)-N-C₁₋₆ alkylamino; C₆₋₁₄ arylamino-carbonylamino which may have substituents; C₆₋₁₄

WO 01/21577

PCT/JP00/06375

7

- arylsulfonylamino which may have substituents; C₆₋₁₄
aryl-carboxyloxy which may have substituents; oxo;
carboxy-C₁₋₆ alkyl; C₁₋₆ alkoxy-carbonyl-C₁₋₆ alkyl; C₇₋₁₉
aralkyl which may have substituents; aromatic hetero
5 ring-C₁₋₆ alkoxy; and cyano;
(8) an antagonist according to the above (1), wherein Ar¹
is piperidinyl, piperazinyl, pyrrolidinyl, dihydropyridyl
or tetrahydropyridyl; each of which may have 1 or 2
substituents selected from the group consisting of oxo, C₆₋₁₄
10 aryl which may have substituents, hydroxy, C₇₋₁₉
aralkyloxy-carbonyl, and C₇₋₁₉ aralkyl;
(9) an antagonist according to the above (1), wherein the
"spacer having a main chain of 1 to 6 atoms" for X and Y
is a bivalent group consisting of 1 to 3 species selected
15 from -O-, -S-, -CO-, -SO-, -SO₂-, -NR⁸- (R⁸ is hydrogen
atom, optionally halogenated C₁₋₆ alkyl, optionally
halogenated C₁₋₆ alkyl-carbonyl, optionally halogenated C₁₋₆
alkylsulfonyl), and a bivalent C₁₋₆ non-cyclic hydrocarbon
group which may have substituents;
20 (10) an antagonist according to the above (1), wherein X
is -CONR^{8c}-, -NR^{8c}CO-, -CH=CH-CONR^{8c}- or -SO₂NR^{8c}-
wherein R^{8c} is hydrogen atom or C₁₋₆ alkyl;
(11) an antagonist according to the above (1), wherein Y
is an optionally halogenated bivalent C₁₋₆ non-cyclic
25 hydrocarbon group;
(12) an antagonist according to the above (1), wherein Ar
is a ring of the formula :



- wherein ----- is a single bond or double bond, n is an integer
30 of 1 to 4;
(13) an antagonist according to the above (1), wherein R¹
and R² are hydrogen atom or C₁₋₆ alkyl which may have
substituents; or R¹ and R², together with the adjacent
nitrogen atom, form a 3 to 8 membered nitrogen-containing

WO 01/21577

PCT/JP00/06375

8

hetero ring;


(14) an antagonist according to the above (1), which is an agent for preventing or treating diseases caused by a melanin-concentrating hormone;

5 (15) an antagonist according to the above (1), which is an agent for preventing or treating obesity;


(16) an antagonist according to the above (1), which is an anorectic agent;

(17) a pharmaceutical, which comprises a melanin-concentrating hormone antagonist in combination with at least one species selected from the group consisting of an agent for treating diabetes, an agent for treating hypertension and an agent for treating arteriosclerosis;

(18) a compound of the formula :



15 wherein Ar¹ is a cyclic group which may have substituents; Ar' is a ring of the formula :



20 wherein ----- is a single bond or double bond, n is an integer of 1 to 4, and each ring may have substituents;

X' is -CONR^{8c}-, -NR^{8c}CO-, -CH=CH-CONR^{8c}- or -SO₂NR^{8c}- where R^{8c} is hydrogen atom or C₁₋₆ alkyl;

Y is a spacer having a main chain of 1 to 6 atoms;


R¹ and R² are independently hydrogen atom or a hydrocarbon group which may have substituents; R¹ and R², together with the adjacent nitrogen atom, may form a nitrogen-containing hetero ring which may have substituents; or R², together with the adjacent nitrogen atom and Y, may form a nitrogen-containing hetero ring which may have substituents;

30 provided that Ar' is a ring of the formula :


WO 01/21577

PCT/JP00/06375


9



wherein symbols have the same meanings as defined above,
and each ring may have substituents, when X' is -SO₂NH-;
and provided that Ar¹ is not biphenyl which may be
substituted, when X' is -CONH- and Ar' is any one of
5 benzopyran, dihydrobenzopyran, dihydrobenzoxazine,
dihydrobenzoxazole or tetrahydrobenzoxazepine;
(excluding N-[2-(N,N-dimethylamino)methyl-6-
tetralinyl]-4-biphenylcarboxamide); or a salt thereof;
10 (19) a compound of the formula :



wherein Ar¹ is a cyclic group which may have substituents;
----- is a single bond or double bond;
n is an integer of 1 to 4;
15 X' is -CONR^{8c}-, -NR^{8c}CO- or -CH=CH-CONR^{8c}- where R^{8c} is
hydrogen atom or C₁₋₆ alkyl;
Y is a spacer having a main chain of 1 to 6 atoms;
R¹ and R² are independently hydrogen atom or a hydrocarbon
group which may have substituents; R¹ and R², together with
20 the adjacent nitrogen atom, may form a nitrogen-containing
hetero ring which may have substituents; or R², together
with the adjacent nitrogen atom and Y, may form a
nitrogen-containing hetero ring which may have
substituents;
25 a ring of the formula :



wherein symbols have the same meanings as defined above,


WO 01/21577

PCT/JP00/06375

10

may have further substituents;
provided that N-[2-(N,N-dimethylamino)methyl-6-tetralinyl]-4-biphenylcarboxamide is excluded; or a salt thereof;


5 (20) a compound according to the above (19), which is of the formula :



wherein R¹ and R² are independently hydrogen atom or a hydrocarbon group which may have substituents; R¹ and R², together with the adjacent nitrogen atom, may form a nitrogen-containing hetero ring which may have substituents; the other symbols have the same meanings as defined in the above (19);

(21) a compound according to the above (20), wherein Ar¹ is an aromatic group which may have substituents; and "a hydrocarbon group which may have substituents" for R¹ and R² is "C₁₋₆ alkyl which may have substituents";

(22) a compound of the formula :



20 wherein Ar¹ is a cyclic group which may have substituents; n is an integer of 1 to 4;

X' is -CONR^{8c}-, -NR^{8c}CO- or -CH=CH-CONR^{8c}- where R^{8c} is hydrogen atom or C₁₋₆ alkyl;

Y is a spacer having a main chain of 1 to 6 atoms;


25 R¹ and R² are independently hydrogen atom or a hydrocarbon group which may have substituents; R¹ and R², together with the adjacent nitrogen atom, may form a nitrogen-containing hetero ring which may have substituents; or R², together with the adjacent nitrogen atom and Y, may form a
30 nitrogen-containing hetero ring which may have

WO 01/21577


PCT/JP00/06375

11


substituents;
a ring of the formula :



wherein n has the same meaning as defined above, may have
5 further substituents;
provided that N-[2-(N,N-dimethylamino)methyl-6-
tetralinyl]-4-biphenylcarboxamide is excluded; or a salt
thereof;
(23) a compound according to the above (22), which is of
10 the formula :



wherein R¹ and R² are independently hydrogen atom or a
hydrocarbon group which may have substituents; R¹ and R²,
together with the adjacent nitrogen atom, may form a
15 nitrogen-containing hetero ring which may have
substituents; the other symbols have the same meanings as
defined in the above (22);
(24) a compound according to the above (23), wherein Ar¹
is an aromatic group which may have substituents; and "a
20 hydrocarbon group which may have substituents" for R¹ and
R² is "C₁-₆ alkyl which may have substituents";
(25) a compound of the formula :



wherein Ar¹ is a cyclic group which may have substituents;
25 X' is -CONR^{8c}-, -NR^{8c}CO- or -CH=CH-CONR^{8c}- where R^{8c} is
hydrogen atom or C₁-₆ alkyl;
Y is a spacer having a main chain of 1 to 6 atoms;
R¹ and R² are independently hydrogen atom or a hydrocarbon


WO 01/21577

PCT/JP00/06375

12


group which may have substituents; R^1 and R^2 , together with the adjacent nitrogen atom, may form a nitrogen-containing hetero ring which may have substituents; or R^2 , together with the adjacent nitrogen atom and Y, may form a nitrogen-containing hetero ring which may have substituents;

a ring of the formula :



may have further substituents; or a salt thereof;


(26) a compound according to the above (25), which is of the formula :



wherein R^1 and R^2 are independently hydrogen atom or a hydrocarbon group which may have substituents; R^1 and R^2 , together with the adjacent nitrogen atom, may form a nitrogen-containing hetero ring which may have substituents; the other symbols have the same meanings as defined in the above (25);

(27) a compound according to the above (26), wherein Ar^1 is an aromatic group which may have substituents; and "a hydrocarbon group which may have substituents" for R^1 and R^2 is " C_{1-6} alkyl which may have substituents";

(28) a compound of the formula :



wherein Ar^1 is a cyclic group which may have substituents; X' is $-CONR^{8c}-$, $-NR^{8c}CO-$, $-CH=CH-CONR^{8c}-$ or $-SO_2NR^{8c}-$ where R^{8c} is hydrogen atom or C_{1-6} alkyl;

Y is a spacer having a main chain of 1 to 6 atoms;


WO 01/21577

PCT/JP00/06375


13

R^1 and R^2 are independently hydrogen atom or a hydrocarbon group which may have substituents; R^1 and R^2 , together with the adjacent nitrogen atom, may form a nitrogen-containing hetero ring which may have substituents; or R^2 , together with the adjacent nitrogen atom and Y, may form a nitrogen-containing hetero ring which may have substituents;

a ring of the formula :



may have further substituents; provided that Ar^1 is not biphenyl which may be substituted, when X' is $-CONH-$; or a salt thereof; (29) a compound of the formula :




wherein Ar^1 is a cyclic group which may have substituents; X' is $-CONR^{8c}-$, $-NR^{8c}CO-$, $-CH=CH-CONR^{8c}-$ or $-SO_2NR^{8c}-$ where R^{8c} is hydrogen atom or C_{1-6} alkyl;

Y is a spacer having a main chain of 1 to 6 atoms;

R^1 and R^2 are independently hydrogen atom or a hydrocarbon group which may have substituents; R^1 and R^2 , together with the adjacent nitrogen atom, may form a nitrogen-containing hetero ring which may have substituents; or R^2 , together with the adjacent nitrogen atom and Y, may form a nitrogen-containing hetero ring which may have substituents;

a ring of the formula :




may have further substituents; or a salt thereof;

(30) a compound of the formula :

WO 01/21577

PCT/JP00/06375


14




wherein Ar¹ is a cyclic group which may have substituents;
X' is -CONR^{8c}-, -NR^{8c}CO-, -CH=CH-CONR^{8c}- or -SO₂NR^{8c}- where
R^{8c} is hydrogen atom or C₁₋₆ alkyl;

- 5 Y is a spacer having a main chain of 1 to 6 atoms;
R¹ and R² are independently hydrogen atom or a hydrocarbon
group which may have substituents; R¹ and R², together with
the adjacent nitrogen atom, may form a nitrogen-containing
hetero ring which may have substituents; or R², together
10 with the adjacent nitrogen atom and Y, may form a
nitrogen-containing hetero ring which may have
substituents;

a ring of the formula :



- 15 may have further substituents; or a salt thereof;
(31) a compound of the formula :



wherein Ar¹ is a cyclic group which may have substituents;
X' is -CONR^{8c}-, -NR^{8c}CO-, -CH=CH-CONR^{8c}- or -SO₂NR^{8c}- where
20 R^{8c} is hydrogen atom or C₁₋₆ alkyl;


- Y is a spacer having a main chain of 1 to 6 atoms;
R¹ and R² are independently hydrogen atom or a hydrocarbon
group which may have substituents; R¹ and R², together with
the adjacent nitrogen atom, may form a nitrogen-containing
hetero ring which may have substituents; or R², together
25 with the adjacent nitrogen atom and Y, may form a
nitrogen-containing hetero ring which may have
substituents;

WO 01/21577

PCT/JP00/06375

15

a ring of the formula :



may have further substituents;

provided that Ar¹ is not biphenyl which may be

5 substituted, when X' is -CONH-; or a salt thereof;

(32) a pharmaceutical composition which comprises a compound as defined in any one of the above (18), (19), (22), (25), (26), (28), (29), (30) and (31);

(33) a prodrug of a compound as defined in any one of the
10 above (18), (19), (22), (25), (26), (28), (29), (30) and (31);

(34) a compound according to the above (18), which is N-[2-(N,N-dimethylamino)methyl-6-tetralinyl]-(4'-methoxybiphenyl-4-yl)carboxamide;

15 4'-fluoro-N-[6-[(N,N-dimethylamino)methyl]-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide;

4'-fluoro-N-[6-(1-piperidinylmethyl)-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide;

20 4'-fluoro-N-[6-[(N,N-dimethylamino)methyl]-5,6,7,8-tetrahydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide;

(+)-4'-fluoro-N-[6-[(N,N-dimethylamino)methyl]-5,6,7,8-tetrahydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide;

25 (-)-4'-fluoro-N-[6-[(N,N-dimethylamino)methyl]-5,6,7,8-tetrahydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide;

4'-chloro-N-[3-[(N,N-dimethylamino)methyl]-2H-chromen-7-yl][1,1'-biphenyl]-4-carboxamide;

30 4'-fluoro-N-[6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide;

N-[3-[(dimethylamino)methyl]-2H-chromen-7-yl]-4'-fluoro[1,1'-biphenyl]-4-carboxamide;

WO 01/21577

PCT/JP00/06375

16

- 4'-chloro-N-[6-[(dimethylamino)methyl]-5-methyl-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide;
6-(4-methoxyphenyl)-N-[5-methyl-6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl]nicotinamide;
5 4'-chloro-N-[7-[(dimethylamino)methyl]-5,6-dihydro-3-quinolinyl][1,1'-biphenyl]-4-carboxamide;
4-(4-chlorophenyl)-N-[6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl]-3,6-dihydro-1(2H)-pyridinecarboxamide;
10 N-[6-[(dimethylamino)methyl]-7,8-dihydro-2-naphthalenyl]-4-(4-fluorophenyl)-1-piperidinecarboxamide;
4-(4-methoxyphenyl)-N-[6-(1-pyrrolidinylmethyl)-5-methyl-7,8-dihydro-2-naphthalenyl]-1-piperidinecarboxamide;
15 4'-fluoro-N-[6-[2-(1-pyrrolidinyl)ethyl]-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide;
4'-chloro-N-[6-[2-(1-pyrrolidinyl)ethyl]-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide;
20 4'-chloro-N-[2-[(dimethylamino)methyl]-3,4-dihydro-2H-1,4-benzoxazin-6-yl][1,1'-biphenyl]-4-carboxamide;
4-(4-methoxyphenyl)-N-[5-methyl-6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl]-1-piperidinecarboxamide;
25 4-(4-chlorophenyl)-N-[6-[(4-methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenyl]-1-piperidinecarboxamide;
4'-chloro-N-[2-[(dimethylamino)methyl]-1H-inden-6-yl][1,1'-biphenyl]-4-carboxamide;
30 4'-fluoro-N-[2-(1-pyrrolidinylmethyl)-3,4-dihydro-2H-1,4-benzoxazin-6-yl][1,1'-biphenyl]-4-carboxamide;
4'-fluoro-N-[5-methyl-6-[(4-methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide;
35 4'-chloro-N-[5-methyl-6-[(4-methyl-1-

- piperaziny]methyl]-7,8-dihydro-2-naphthalenyl][1,1'-
biphenyl]-4-carboxamide; or
4-(4-chlorophenyl)-N-[5-methyl-6-[(4-methyl-1-
piperaziny]methyl]-7,8-dihydro-2-naphthalenyl]-1-
5 piperidinecarboxamide;
(35) a method for preventing or treating diseases caused
by a melanin-concentrating hormone in a mammal in need
thereof, which comprises administering to said mammal an
effective amount of a compound or a salt thereof as defined
10 in the above (1);
(36) a method for preventing or treating obesity in a mammal
in need thereof, which comprises administering to said
mammal an effective amount of a compound or a salt thereof
as defined in the above (1);
15 (37) use of a compound or a salt thereof as defined in the
above (1), for the manufacture of a pharmaceutical
preparation for preventing or treating diseases caused by
a melanin-concentrating hormone; and
(38) use of a compound or a salt thereof as defined in the
20 above (1), for the manufacture of a pharmaceutical
preparation for preventing or treating obesity.

Examples of "cyclic group" in the "cyclic group which
may have substituents" for Ar¹ include aromatic groups,
25 non-aromatic cyclic hydrocarbon groups, non-aromatic
heterocyclic groups.

Here, examples of "aromatic groups" include
monocyclic aromatic groups, condensed aromatic groups, and
ring assembly aromatic groups.

30 Examples of the concerned monocyclic aromatic groups
include univalent groups which can be formed by removing
an optional one hydrogen atom from a monocyclic aromatic
ring. Example of the "monocyclic aromatic ring" include
a benzene ring and a 5 or 6 membered aromatic hetero ring.

35 Examples of the "5 or 6 membered aromatic hetero ring"
include a 5 or 6 membered aromatic hetero ring containing

WO 01/21577

PCT/JP00/06375

18

one or more (for example, 1 to 3) hetero atom selected from nitrogen, sulfur and oxygen atom in addition to a carbon atom. Concretely, thiophene, furan, pyrrole, imidazole, pyrazole, thiazole, isothiazole, oxazole, isoxazole,
5 pyridine, pyrazine, pyrimidine, pyridazine, 1,2,4-oxadiazole, 1,3,4-oxadiazole, 1,2,4-thiadiazole, 1,3,4-thiadiazole, furazan, etc., can be mentioned.

Concrete examples of the "monocyclic aromatic groups" include phenyl, 2- or 3-thienyl, 2-, 3-, or 4-pyridyl, 2-
10 or 3-furyl, 2-, 4- or 5-thiazonyl, 2-, 4- or 5-oxazolyl, 1-, 3- or 4-pyrazolyl, 2-pyrazinyl, 2-, 4- or 5-pyrimidinyl, 1-, 2- or 3-pyrrolyl, 1-, 2- or 4-imidazolyl, 3- or 4-pyridazinyl, 3-isothiazolyl, 3-isooxazolyl, 1,2,4-oxadiazol-5-yl, 1,2,4-oxadiazol-3-yl.

15 The "condensed aromatic groups" mean a univalent group that can be formed by removing an optional one hydrogen atom from condensed polycyclic (preferably bicyclic to tetracyclic, more preferably bicyclic or tricyclic) aromatic rings. Examples of the "condensed aromatic
20 groups" include condensed polycyclic aromatic hydrocarbons, condensed polycyclic aromatic hetero rings.

Examples of the "condensed polycyclic aromatic hydrocarbons" include C₉₋₁₄ condensed polycyclic (bicyclic or tricyclic) aromatic hydrocarbons (e.g. naphthalene,
25 indene, fluorene, anthracene, etc.).

Examples of the "condensed polycyclic aromatic hetero rings" include 9 to 14 membered, preferably, 9 or 10 membered, condensed polycyclic aromatic hetero rings containing one or more (for instance, 1 to 4 atoms) hetero
30 atoms selected from nitrogen, sulfur and oxygen atom in addition to carbon atoms. Concrete examples of the "condensed polycyclic aromatic hetero rings" include benzofuran, benzimidazole, benzoxazole, benzothiazole, benzisothiazole, naphtho[2,3-b]thiophene, isoquinoline, quinoline, indole, quinoxaline, phenanthridine,
35 phenothiadine, phenoxazine, phthaladine, naphthylidine,

quinazoline, cinnoline, carbazole, β -carboline, acridine, phenazine, phthalimide, thioxanthene.

Concrete examples of "condensed aromatic groups" include 1-naphthyl; 2-naphthyl; 2-, 3-, 4-, 5- or 8-quinolyl; 1-, 3-, 4-, 5-, 6-, 7- or 8-isoquinolyl; 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl; 1-, 2-, 4- or 5-isoindolyl; 1-, 5- or 6-phthalazinyl; 2-, 3- or 5-quinoxalyl; 2-, 3-, 4-, 5- or 6-benzofuranyl; 2-, 4-, 5- or 6-benzothiazolyl; 1-, 2-, 4-, 5- or 6-benzimidazolyl; 1-, 2-, 3- or 4-fluorenyl; thioxanthenyl.

"Ring assembly aromatic group" means a group formed by removing an optional one hydrogen atom from an aromatic ring assemblies in which 2 or more (preferably 2 or 3) aromatic rings are directly bonded by single bonds, and in which the number of bonds which directly bond the rings, is less by one than the number of ring systems.

Examples of the aromatic ring assemblies include an aromatic ring assemblies formed by 2 or 3 (preferably 2) species selected from C_{6-14} monocyclic or condensed polycyclic aromatic hydrocarbons (e.g. benzene and naphthalene) and 5 to 10 membered (preferably 5 or 6 membered) aromatic hetero rings.

Preferable example of the aromatic ring assemblies include aromatic ring assemblies comprising 2 or 3 aromatic rings selected from benzene, naphthalene, pyridine, pyrimidine, thiophene, furan, thiazole, isothiazole, oxazole, isoxazole, 1,2,4-oxadiazole, 1,3,4-oxadiazole, 1,2,4-thiadiazole, 1,3,4-thiadiazole, quinoline, isoquinoline, indole, benzothiophene, benzoxazole, benzothiazole, benzofuran and pyrrole.

Concrete examples of the "ring assembly aromatic groups" include 2-, 3- or 4-biphenyl; 3-(1-naphthyl)-1,2,4-oxadiazol-5-yl; 3-(2-naphthyl)-1, 2, 4-oxadiazol-5-yl; 3-(2-benzofuranyl)-1,2,4-oxadiazol-5-yl; 3-phenyl-1,2,4-oxadiazol-5-yl; 3-(2-benzoxazolyl)-1,2,4-oxadiazol-5-yl; 3-(3-indolyl)-1,2,4-oxadiazol-5-yl; 3-

WO 01/21577

PCT/JP00/06375

20

(2-indolyl)-1,2,4-oxadiazol-5-yl; 4-phenylthiazol-2-yl;
4-(2-benzofuranyl)thiazol-2-yl; 4-phenyl-1,3-oxazol-5-
yl; 5-phenyl-isothiazol-4-yl; 5-phenyloxazol-2-yl; 4-
(2-thienyl)phenyl; 4-(3-thienyl)phenyl; 3-(3-
5 pyridyl)phenyl; 4-(3-pyridyl)phenyl; 6-phenyl-3-pyridyl;
5-phenyl-1,3,4-oxadiazol-2-yl; 4-(2-naphthyl)phenyl; 4-
(2-benzofuranyl)phenyl; 4,4'-terphenyl; 5-phenyl-2-
pyridyl; 2-phenyl-5-pyrimidinyl; 4-(4-pyridyl)phenyl;
2-phenyl-1,3-oxazol-5-yl; 2,4-diphenyl-1,3-oxazol-5-yl;
10 3-phenyl-isoxazol-5-yl; 5-phenyl-2-furyl; 4-(2-
furyl)phenyl; 3-(4-pyridyl)pyrrolyl.

Preferable groups among the above "aromatic groups"
are "C₆₋₁₄ monocyclic or condensed polycyclic aromatic
hydrocarbon groups (preferably, phenyl, etc.)", "a group
15 formed by removing an optional one hydrogen atom from an
aromatic ring assemblies in which 2 or 3 C₆₋₁₄ monocyclic or
condensed polycyclic aromatic hydrocarbon groups are
directly bonded by single bonds (preferably, 2-, 3- or
4-biphenyl; 4,4'-terphenyl, etc.)" and "a group formed by
20 removing an optional one hydrogen atom from an aromatic ring
assemblies in which a C₆₋₁₄ monocyclic or condensed polycyclic
aromatic hydrocarbon and 5 to 10 membered aromatic hetero
ring are directly bonded by a single bond (preferably,
6-phenyl-3-pyridyl, 5-phenyl-2-pyridyl, etc.)".

25 Examples of "non-aromatic cyclic hydrocarbon groups"
include C₃₋₈ Cycloalkyl, C₃₋₈ cycloalkenyl.

Here, concrete examples of C₃₋₈ cycloalkyl include
cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl,
cycloheptyl, cyclooctyl.

30 Concrete examples of C₃₋₈ cycloalkenyl include
cyclopropenyl, cyclobutenyl, cyclopentenyl,
cyclohexenyl, cycloheptenyl, cyclooctenyl.

Among the above "non-aromatic cyclic hydrocarbon
groups", C₃₋₈ cycloalkyl is preferable, and cyclohexyl is
35 particularly preferable.

Examples of "non-aromatic heterocyclic groups"

WO 01/21577

PCT/JP00/06375

21

include monocyclic non-aromatic heterocyclic groups,
condensed polycyclic non-aromatic heterocyclic groups.

Examples of the "monocyclic non-aromatic heterocyclic groups" include univalent groups formed by removing an optional one hydrogen atom from monocyclic non-aromatic hetero ring. Examples of the "monocyclic non-aromatic heterocyclic groups" include 5 to 8 membered monocyclic non-aromatic heterocyclic groups containing one or more (e.g. 1 to 3) hetero atoms selected from nitrogen, sulfur and oxygen atom in addition to carbon atoms. Concretely, tetrahydrothiophene, tetrahydrofuran, pyrrolidine, imidazoline, imidazolidine, pyrazoline, pyrazolidine, tetrahydrothiazole, tetrahydroisothiazole, tetrahydrooxazole, tetrahydroisoxazole, piperidine, tetrahydropyridine, dihydropyridine, piperazine, morpholine, thiomorpholine, tetrahydropyrimidine, tetrahydropyridazine, hexamethyleneimine, etc. can be mentioned.

"Condensed polycyclic non-aromatic heterocyclic group" means a univalent group formed by removing an optional one hydrogen atom from a condensed polycyclic (preferably bicyclic to tetracyclic, more preferably bicyclic or tricyclic) non-aromatic hetero ring. Examples of the "condensed polycyclic non-aromatic hetero ring" include 9 to 14 membered, preferably 9 or 10 membered condensed polycyclic non-aromatic hetero rings which contain one or more (e.g. 1 to 4) hetero atoms selected from nitrogen, sulfur and oxygen atom in addition to carbon atoms.

Concretely, dihydrobenzofuran, dihydrobenzimidazole, dihydrobenzoxazole, dihydrobenzothiazole, dihydrobenzisothiazole, dihydronaphtho[2,3-b]thiophene, tetrahydroisoquinoline, tetrahydroquinoline, indoline, isoindoline, tetrahydroquinoxaline, tetrahydrophenanthridine, hexahydrophenothiadine, hexahydrophenoxazine,

tetrahydrophthaladine, tetrahydronaphthylidine,
tetrahydroquinazoline, tetrahydrocinnoline,
tetrahydrocarbazole, tetrahydro- β -carboline,
tetrahydroacridine, tetrahydrophenazine,
5 tetrahydrothioxantene, etc., can be mentioned.

Among the above "non-aromatic heterocyclic groups",
"5 to 8 membered monocyclic non-aromatic heterocyclic
groups (preferably piperidinyl; piperazinyl;
pyrrolidinyl; dihydropyridyl; tetrahydropyridyl, etc.)"
10 are preferable.

Examples of "substituents" in the "cyclic group which
may have substituents" for Ar¹ include oxo, halogen atoms
(e.g. fluorine, chlorine, bromine, iodine, etc.), C₁₋₃
15 alkylenedioxy (e.g. methylenedioxy, ethylenedioxy, etc.),
nitro, cyano, optionally halogenated C₁₋₆ alkyl, hydroxy-C₁₋₆
alkyl, carboxy-C₁₋₆ alkyl, C₁₋₆ alkoxy-carbonyl-C₁₋₆ alkyl,
C₆₋₁₄ aryloxy-C₁₋₆ alkyl (e.g. phenoxymethyl, etc.), C₁₋₆
alkyl-C₆₋₁₄ aryl-C₂₋₆ alkenyl (e.g. methylphenylethenyl,
20 etc.), optionally halogenated C₃₋₆ cycloalkyl, optionally
halogenated C₁₋₆ alkoxy, optionally halogenated C₁₋₆
alkylthio, C₇₋₁₉ aralkyl which may have substituents,
hydroxy, C₆₋₁₄ aryloxy which may have substituents, C₇₋₁₉
aralkyloxy which may have substituents, C₆₋₁₄ aryl-carbamoyl
25 which may have substituents, amino, amino-C₁₋₆ alkyl (e.g.
aminomethyl, aminoethyl, aminopropyl, aminobutyl, etc.),
mono-C₁₋₆ alkylamino (e.g. methylamino, ethylamino,
propylamino, isopropylamino, butylamino, etc.), di-C₁₋₆
alkylamino (e.g. dimethylamino, diethylamino,
30 dipropylamino, dibutylamino, ethylmethylamino, etc.),
mono-C₁₋₆ alkylamino-C₁₋₆ alkyl (e.g. methylaminomethyl,
ethylaminomethyl, propylaminomethyl,
isopropylaminoethyl, butylaminoethyl, etc.), di-C₁₋₆
alkylamino-C₁₋₆ alkyl (e.g. dimethylaminomethyl,
35 diethylaminomethyl, dipropylaminomethyl,
diisopropylaminoethyl, dibutylaminoethyl, etc.), 5 to 7

WO 01/21577

PCT/JP00/06375

23

membered saturated cyclic amino which may have substituents, 5 to 7 membered non-aromatic heterocyclic groups which may have substituents, acyl, acylamino, acyloxy, aromatic hetero ring-C₁₋₆ alkoxy.

5 The "cyclic group" for Ar¹ may have 1 to 5, preferably 1 to 3, of the above-mentioned substituents at a substitutable position on the cyclic group. When the number of substituents is 2 or more, each substituents can be the same or different.

10 Also, when the "cyclic group" for Ar¹ is a non-aromatic cyclic hydrocarbon group or a non-aromatic heterocyclic group, the "cyclic group" may have as its substituents, C₆₋₁₄ aryl which may have substituents, and 5 to 10 membered aromatic heterocyclic groups which may have substituents.

15 Here, the groups exemplified as "substituents" in the "5 to 7 membered saturated cyclic amino which may have substituents" mentioned hereinafter, can be mentioned as "C₆₋₁₄ aryl which may have substituents" and "5 to 10 membered aromatic heterocyclic groups which may have substituents".

20 The number of substituents is, for instance, 1 to 3. When the number of substituents is 2 or more, each substituents can be the same or different.

Concrete examples of the above "optionally halogenated C₁₋₆ alkyl" include C₁₋₆ alkyl (e.g. methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, hexyl, etc.) which may have 1 to 5, preferably 1 to 3, halogen atoms (e.g. fluorine, chlorine, bromine, iodine, etc.). Concrete examples include methyl, chloromethyl, difluoromethyl, trichloromethyl, trifluoromethyl, ethyl, 2-bromoethyl, 2,2,2-trifluoroethyl, pentafluoroethyl, propyl, 3,3,3-trifluoropropyl, isopropyl, butyl, 4,4,4-trifluorobutyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, neopentyl, 5,5,5-trifluoropentyl, hexyl, 6,6,6-trifluorohexyl.

The C₁₋₆ alkyl in the above "optionally halogenated C₁₋₆

alkyl" can be mentioned as the C₁₋₆ alkyl in the above
"hydroxy-C₁₋₆ alkyl", "carboxy-C₁₋₆ alkyl" and "C₁₋₆
alkoxy-carbonyl-C₁₋₆ alkyl". Examples of C₁₋₆ alkoxy in the
"C₁₋₆ alkoxy-carbonyl-C₁₋₆ alkyl" include methoxy, ethoxy,
5 propoxy, butoxy, pentyloxy.

Examples of the above "optionally halogenated C₃₋₆
cycloalkyl" include C₃₋₆ cycloalkyl (e.g. cyclopropyl,
cyclobutyl, cyclopentyl, cyclohexyl, etc.) which may have
1 to 5, preferably 1 to 3, halogen atoms (e.g. fluorine,
10 chlorine, bromine, iodine, etc.). Concrete examples
include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl,
4,4-dichlorocyclohexyl, 2,2,3,3-tetrafluorocyclopentyl,
4-chlorocyclohexyl.

Examples of the above "optionally halogenated C₁₋₆
15 alkoxy" include C₁₋₆ alkoxy (e.g. methoxy, ethoxy, propoxy,
butoxy, pentyloxy, etc.) which may have 1 to 5, preferably
1 to 3, halogen atoms (e.g. fluorine, chlorine, bromine,
iodine, etc.). Concrete examples include methoxy,
difluoromethoxy, trifluoromethoxy, ethoxy, 2,2,2-
20 trifluoroethoxy, propoxy, isopropoxy, butoxy, 4,4,4-
trifluorobutoxy, isobutoxy, sec-butoxy, pentyloxy,
hexyloxy.

Examples of the above "optionally halogenated C₁₋₆
alkylthio" include C₁₋₆ alkylthio (e.g. methylthio,
25 ethylthio, propylthio, isopropylthio, butylthio, sec-
butylthio, tert-butylthio, etc.) which may have 1 to 5,
preferably 1 to 3, halogen atoms (e.g. fluorine, chlorine,
bromine, iodine, etc.). Concrete examples include
methylthio, difluoromethylthio, trifluoromethylthio,
30 ethylthio, propylthio, isopropylthio, butylthio, 4,4,4-
trifluorobutylthio, pentylthio, hexylthio.

Examples of the "C₇₋₁₉ aralkyl" in the above "C₇₋₁₉
aralkyl which may have substituents" include benzyl,
phenethyl, diphenylmethyl, triphenylmethyl, 1-
35 naphthylmethyl, 2-naphthylmethyl, 2,2-diphenylethyl, 3-
phenylpropyl, 4-phenylbutyl, 5-phenylpentyl. Benzyl is

particularly preferable.

Examples of the "substituents" in the above "C₇₋₁₉ aralkyl which may have substituents" include halogen atom (e.g. fluorine, chlorine, bromine, iodine, etc.), C₁₋₃ alkylene dioxy (e.g. methylenedioxy, ethylenedioxy, etc.), nitro, cyano, optionally halogenated C₁₋₆ alkyl, optionally halogenated C₃₋₆ cycloalkyl, optionally halogenated C₁₋₆ alkoxy, optionally halogenated C₁₋₆ alkylthio, hydroxy, amino, mono-C₁₋₆ alkylamino (e.g. methylamino, ethylamino, propylamino, isopropylamino, butylamino, etc.), di-C₁₋₆ alkylamino (e.g. dimethylamino, diethylamino, dipropylamino, dibutylamino, ethylmethylamino, etc.), amino-C₁₋₆ alkyl (e.g. aminomethyl, aminoethyl, aminopropyl, aminobutyl, etc.), mono-C₁₋₆ alkylamino-C₁₋₆ alkyl (e.g. methylaminomethyl, ethylaminomethyl, propylaminomethyl, isopropylaminoethyl, butylaminoethyl, etc.), di-C₁₋₆ alkylamino-C₁₋₆ alkyl (e.g. dimethylaminomethyl, diethylaminomethyl, dipropylaminomethyl, diisopropylaminoethyl, dibutylaminoethyl, etc.), formyl, carboxy, carbamoyl, thiocarbamoyl, optionally halogenated C₁₋₆ alkyl-carbonyl, C₁₋₆ alkoxy-carbonyl (e.g., methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, tert-butoxycarbonyl, etc.), mono-C₁₋₆ alkyl-carbamoyl (e.g., methylcarbamoyl, ethylcarbamoyl, etc.), di-C₁₋₆ alkyl-carbamoyl (e.g. dimethylcarbamoyl, diethylcarbamoyl, ethylmethylcarbamoyl, etc.), optionally halogenated C₁₋₆ alkylsulfonyl, formylamino, optionally halogenated C₁₋₆ alkyl-carboxamide, C₁₋₆ alkoxy-carboxamide (e.g. methoxycarboxamide, ethoxycarboxamide, propoxycarboxamide, butoxycarboxamide, etc.), C₁₋₆ alkylsulfonylamino (e.g. methylsulfonylamino, ethylsulfonylamino, etc.), C₁₋₆ alkyl-carbonyloxy (e.g. acetoxyl, propanoyloxy, etc.), C₁₋₆ alkoxy-carbonyloxy (e.g. methoxycarbonyloxy, ethoxycarbonyloxy, propoxycarbonyloxy, butoxycarbonyloxy, etc.) mono-C₁₋₆

alkyl-carbamoyloxy (e.g. methylcarbamoyloxy, ethylcarbamoyloxy, etc.), di-C₁₋₆ alkyl-carbamoyloxy (e.g. dimethylcarbamoyloxy, diethylcarbamoyloxy, etc.). The number of substituents is, for instance, 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituents can be the same or different.

As "optionally halogenated C₁₋₆ alkyl", "optionally halogenated C₃₋₆ cycloalkyl", "optionally halogenated C₁₋₆ alkoxy" and "optionally halogenated C₁₋₆ alkylthio", those exemplified as "substituents" in the above "cyclic group which may have substituents" can be used respectively.

Examples of the above "optionally halogenated C₁₋₆ alkylcarbonyl" include C₁₋₆ alkyl-carbonyl (e.g. acetyl, propanoyl, butanoyl, pentanoyl, hexanoyl, etc.) which may have 1 to 5, preferably 1 to 3, halogen atoms (e.g., fluorine, chlorine, bromine, iodine, etc.). Concrete examples include acetyl, monochloroacetyl, trifluoroacetyl, trichloroacetyl, propanoyl, butanoyl, pentanoyl, hexanoyl.

Examples of the above "optionally halogenated C₁₋₆ alkylsulfonyl" include C₁₋₆ alkylsulfonyl (e.g. methylsulfonyl, ethylsulfonyl, propylsulfonyl, isopropylsulfonyl, butylsulfonyl, sec-butylsulfonyl, tert-butylsulfonyl, etc.) which may have 1 to 5, preferably 1 to 3, halogen atoms (e.g., fluorine, chlorine, bromine, iodine, etc.). Concrete examples include methylsulfonyl, difluoromethylsulfonyl, trifluoromethylsulfonyl, ethylsulfonyl, propylsulfonyl, isopropylsulfonyl, butylsulfonyl, 4,4,4-trifluorobutylsulfonyl, pentylsulfonyl, hexylsulfonyl.

Examples of the above "optionally halogenated C₁₋₆ alkyl-carboxamide" include C₁₋₆ alkyl-carboxamide (e.g. acetamide, propanamide, butanamide, etc.) which may have 1 to 5, preferably 1 to 3, halogen atoms (e.g. fluorine, chlorine, bromine, iodine, etc.). Concrete examples include acetamide, trifluoroacetamide, propanamide,

WO 01/21577

PCT/JP00/06375

27

butanamide.

Examples of "C₆₋₁₄ aryloxy" in the above "C₆₋₁₄ aryloxy which may have substituents" include phenyloxy, 1-naphthyloxy, 2-naphthyloxy.

5 Examples of "C₇₋₁₉ aralkyloxy" in the above "C₇₋₁₉ aralkyloxy which may have substituents" include benzyloxy, phenethyloxy, diphenylmethyloxy, triphenylmethyloxy, 1-naphthylmethyloxy, 2-naphthylmethyloxy, 2,2-diphenylethyloxy, 3-phenylpropyloxy, 4-phenylbutyloxy,
10 5-phenylpentyloxy.

Examples of "C₆₋₁₄ arylcarbamoyl" in the above "C₆₋₁₄ arylcarbamoyl which may have substituents" include phenylcarbamoyl, 1-naphthylcarbamoyl, 2-naphthylcarbamoyl.

15 As the "substituents" in the "C₆₋₁₄ aryloxy which may have substituents", "C₇₋₁₉ aralkyloxy which may have substituents" and "C₆₋₁₄ aryl-carbamoyl which may have substituents", those exemplified for "substituents" in the above "C₇₋₁₉ aralkyl which may have substituents" can be used.

20 The number of substituents is, for instance, 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituents can be the same or different.

Examples of the "5 to 7 membered saturated cyclic amino" in the above "5 to 7 membered saturated cyclic amino which may have substituents" include morpholino, thiomorpholino, piperazin-1-yl, piperidino, pyrrolidin-1-yl. The "5 to 7 membered saturated cyclic amino" can be condensed with a benzene ring.

25 Examples of "substituents" in the "5 to 7 membered saturated cyclic amino which may have substituents" include oxo, optionally halogenated C₁₋₆ alkyl, optionally halogenated C₁₋₆ alkyl-carbonyl, optionally halogenated C₁₋₆ alkylsulfonyl, C₆₋₁₄ aryl which may have substituents, C₇₋₁₉ aralkyl which may have substituents, C₆₋₁₄ aryl-carbonyl
30 which may have substituents, 5 to 10 membered aromatic heterocyclic group which may have substituents, 5 to 8

WO 01/21577

PCT/JP00/06375

28

membered monocyclic non-aromatic heterocyclic group (e.g., piperidino, piperazinyl, pyrrolidinyl, dihydropyridyl, etc.). The number of substituents is, for instance, 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituents can be the same or different.

Here, as "optionally halogenated C₁₋₆ alkyl" and "C₇₋₁₉ aralkyl which may have substituents", those exemplified as "substituents" in the above "cyclic group which may have substituents" can be used.

As "optionally halogenated C₁₋₆ alkyl-carbonyl" and "optionally halogenated C₁₋₆ alkylsulfonyl", those exemplified as "substituents" in the above "C₇₋₁₉ aralkyl which may have substituents" can be used.

Examples of the "C₆₋₁₄ aryl" in the "C₆₋₁₄ aryl which may have substituents" include phenyl, 1-naphthyl, 2-naphthyl, 2-indenyl, 2-anthryl. Phenyl is especially preferable.

As the substituents in the "C₆₋₁₄ aryl which may have substituents", those exemplified as "substituents" in the above "C₇₋₁₉ aralkyl which may have substituents" can be used.

The number of substituents is, for instance, 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituents can be the same or different.

Examples of the "C₆₋₁₄ aryl-carbonyl" in the "C₆₋₁₄ aryl-carbonyl which may have substituents" include benzoyl, 1-naphthoyl, 2-naphthoyl.

As the "substituents" in the "C₆₋₁₄ aryl-carbonyl which may have substituents", those exemplified as "substituents" in the above "C₇₋₁₉ aralkyl which may have substituents" can be used. The number of substituents is, for instance, 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituents can be the same or different.

Examples of "5 to 10 membered aromatic heterocyclic groups" in "5 to 10 membered aromatic heterocyclic groups which may have substituents" include 5 to 10 membered (monocyclic or bicyclic) aromatic heterocyclic groups

containing 1 or 2 kinds of, preferably 1 to 4 hetero atoms selected from nitrogen, sulfur and oxygen atom in addition to carbon atoms. Concrete examples include 2- or 3-thienyl; 2-, 3- or 4-pyridyl; 2- or 3-furyl; 2-, 4- or 5-thiazolyl; 2-, 4- or 5-oxazolyl; 1-, 3- or 4-pyrazolyl; 2-pyrazinyl; 2-, 4- or 5-pyrimidinyl; 1-, 2- or 3-pyrrolyl; 1-, 2- or 4-imidazolyl; 3- or 4-pyridazinyl; 3-isothiazolyl; 3-isoxazolyl; 1,2,4-oxadiazol-5-yl; 1,2,4-oxadiazol-3-yl; 2-, 3-, 4-, 5- or 8-quinolyl; 1-, 3-, 4-, 5-, 6-, 7- or 8-isoquinolyl; 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl; 1-, 2-, 4- or 5-isoindolyl; 1-, 5- or 6-phthalazinyl; 2-, 3- or 5-quinoxalinyll; 2-, 3-, 4-, 5- or 6-benzofuranyl; 2-, 4-, 5- or 6-benzothiazolyl; 1-, 2-, 4-, 5- or 6-benzimidazolyl.

Examples of the "substituents" in the "5 to 10 membered aromatic heterocyclic groups which may have substituents" include halogen atom (e.g. fluorine, chlorine, bromine and iodine, etc.), C₁₋₃ alkylenedioxy (e.g. methylenedioxy, ethylenedioxy, etc.), nitro, cyano, optionally halogenated C₁₋₆ alkyl, C₆₋₁₄ aryloxy-C₁₋₆ alkyl (e.g. phenoxymethyl, etc.), C₁₋₆ alkyl-C₆₋₁₄ aryl-C₂₋₆ alkenyl (e.g. methylphenylethenyl, etc.), optionally halogenated C₃₋₆ cycloalkyl, optionally halogenated C₁₋₆ alkoxy, optionally halogenated C₁₋₆ alkylthio, C₇₋₁₉ aralkyl which may have substituents, hydroxy, C₆₋₁₄ aryloxy which may have substituents, C₇₋₁₉ aralkyloxy which may have substituents, amino, amino-C₁₋₆ alkyl (e.g. aminomethyl, aminoethyl, aminopropyl, aminobutyl, etc.), mono-C₁₋₆ alkylamino (e.g. methylamino, ethylamino, propylamino, isopropylamino, butylamino, etc.), di-C₁₋₆ alkylamino (e.g. dimethylamino, diethylamino, dipropylamino, dibutylamino, ethylmethylamino, etc.), mono-C₁₋₆ alkylamino-C₁₋₆ alkyl (e.g. methylaminomethyl, ethylaminomethyl, propylaminomethyl, isopropylaminoethyl, butylaminoethyl, etc.), di-C₁₋₆ alkylamino-C₁₋₆ alkyl (e.g. dimethylaminomethyl, diethylaminomethyl,

WO 01/21577

PCT/JP00/06375

30

dipropylaminomethyl, diisopropylaminoethyl, dibutylaminoethyl, etc.), 5 to 7 membered saturated cyclic amino, acyl, acylamino, acyloxy. The number of substituents is, for instance, 1 to 5, preferably 1 to 3.

5 When the number of substituents is 2 or more, each substituents can be the same or different.

Here, as "optionally halogenated C₁₋₆ alkyl", "optionally halogenated C₃₋₆ cycloalkyl", "optionally halogenated C₁₋₆ alkoxy", "optionally halogenated C₁₋₆ alkylthio", "C₇₋₁₉ aralkyl which may have substituents", "C₆₋₁₄ aryloxy which may have substituents", "C₇₋₁₉ aralkyloxy which may have substituents", those exemplified as the "substituent" in the above "cyclic group which may have substituents" can be used respectively.

15 As a "5 to 7 membered saturated cyclic amino", those exemplified as "5 to 7 membered saturated cyclic amino" regarding "5 to 7 membered saturated cyclic amino which may have substituents" which is a "substituent" in the above "5 to 7 membered saturated cyclic amino which may have substituents" can be used.

20 Examples of the above "acyl" include acyl of the formulae : -CO-R³, -CO-OR³, -CO-NR³R⁴, -CS-NR³R⁴, -SO₂-R^{3a}, -SO-R^{3a}, -PO(-OR³)-OR⁴ or -PO₂-R^{3a} wherein R³ is (i) hydrogen atom, (ii) a hydrocarbon group which may have substituents, or (iii) a heterocyclic group which may have substituents; R^{3a} is (i) a hydrocarbon group which may have substituents, or (ii) a heterocyclic group which may have substituents; R⁴ is hydrogen atom or C₁₋₆ alkyl; R³ and R^{3a}, together with the adjacent nitrogen atom, can form a nitrogen-containing hetero ring which may have substituents.

35 Examples of the "hydrocarbon group" in "hydrocarbon group which may have substituents" for R³ or R⁴ include straight-chain or cyclic hydrocarbon groups (e.g. alkyl, alkenyl, alkynyl, cycloalkyl, aryl, aralkyl, etc.). Among these, C₁₋₁₉ straight chain or cyclic hydrocarbon groups as

WO 01/21577

PCT/JP00/06375

31

shown below are preferable.

a) C₁₋₆ alkyl (e.g. methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, hexyl, etc.);

5 b) C₂₋₆ alkenyl (e.g., vinyl, allyl, isopropenyl, 2-butenyl, etc.);

c) C₂₋₆ alkynyl (e.g. ethynyl, propargyl, 2-butyne, etc.);

10 d) C₃₋₆ cycloalkyl (e.g. cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, etc.); the C₃₋₆ cycloalkyl can be condensed with one benzene ring;

e) C₆₋₁₄ aryl (e.g. phenyl, 1-naphthyl, 2-naphthyl, 2-indenyl, 2-anthryl, etc.), preferably phenyl;

15 f) C₇₋₁₉ aralkyl (e.g. benzyl, phenethyl, diphenylmethyl, triphenylmethyl, 1-naphthylmethyl, 2-naphthylmethyl, 2,3-diphenylethyl, 3-phenylpropyl, 4-phenylbutyl, 5-phenylpentyl, etc.), preferably benzyl.

The "hydrocarbon groups" are preferably C₁₋₆ alkyl, C₆₋₁₄ aryl, C₇₋₁₉ aralkyl, etc.

20

Examples of the "substituent" in "hydrocarbon groups which may have substituents" include halogen atom (e.g. fluorine, chlorine, bromine, iodine, etc.), C₁₋₃ alkylenedioxy (e.g. methylenedioxy, ethylenedioxy, etc.),
25 nitro, cyano, optionally halogenated C₁₋₆ alkoxy, optionally halogenated C₁₋₆ alkylthio, hydroxy, amino, mono-C₁₋₆ alkylamino (e.g. methylamino, ethylamino, propylamino, isopropylamino, butylamino, etc.), di-C₁₋₆ alkylamino (e.g. dimethylamino, diethylamino,
30 dipropylamino, dibutylamino, ethylmethylamino, etc.), formyl, carboxy, carbamoyl, thiocarbamoyl, optionally halogenated C₁₋₆ alkyl-carbonyl, C₁₋₆ alkoxy-carbonyl (e.g., methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, tert-butoxycarbonyl, etc.), 5 to 10 membered aromatic
35 heterocyclic groups which may have substituents, C₆₋₁₄ aryl-carbonyl which may have substituents, C₆₋₁₄

aryloxy-carbonyl which may have substituents, C₇₋₁₉
 aralkyloxy-carbonyl which may have substituents, 5 to 6
 membered hetero ring-carbonyl which may have substituents,
 mono-C₁₋₆ alkyl-carbamoyl (e.g. methylcarbamoyl,
 5 ethylcarbamoyl, etc.), di-C₁₋₆ alkyl-carbamoyl (e.g.
 dimethylcarbamoyl, diethylcarbamoyl,
 ethylmethylcarbamoyl, etc.), C₆₋₁₄ aryl-carbamoyl which may
 have substituents, 5 to 6 membered hetero ring-carbamoyl
 which may have substituents, optionally halogenated C₁₋₆
 10 alkylsulfonyl, C₆₋₁₄ arylsulfonyl which may have
 substituents, formylamino, C₁₋₆ alkyl-carbonyloxy (e.g.
 acetoxy, propanoyloxy, etc.), C₆₋₁₄ aryl-carbonyloxy which
 may have substituents, C₁₋₆ alkoxy-carbonyloxy (e.g.
 methoxycarbonyloxy, ethoxycarbonyloxy,
 15 propoxycarbonyloxy, butoxycarbonyloxy, etc.), mono-C₁₋₆
 alkyl-carbamoyloxy (e.g. methylcarbamoyloxy,
 ethylcarbamoyloxy, etc.), di-C₁₋₆ alkyl-carbamoyloxy (e.g.
 dimethylcarbamoyloxy, diethylcarbamoyloxy, etc.), C₆₋₁₄
 aryl-carbamoyloxy which may have substituents,
 20 nicotinoyloxy. The number of substituents is, for
 instance, 1 to 5, preferably 1 to 3. When the number of
 substituents is 2 or more, each substituents can be the same
 or different.

Here, as "optionally halogenated C₁₋₆ alkoxy",
 25 "optionally halogenated C₁₋₆ alkylthio" and "C₆₋₁₄ aryl-
 carbamoyl which may have substituents", those exemplified
 as a "substituent" in the above "cyclic group which may have
 substituents" can be used.

As "optionally halogenated C₁₋₆ alkyl-carbonyl" and
 30 "optionally halogenated C₁₋₆ alkylsulfonyl", those
 exemplified as a "substituent" in the above "C₇₋₁₉ aralkyl
 which may have substituents" can be used.

As the above "5 to 10 membered aromatic heterocyclic
 groups which may have substituents" and "C₆₋₁₄ aryl-carbonyl
 35 which may have substituents", those exemplified as
 "substituent" in the above "5 to 7 membered saturated cyclic

WO 01/21577

PCT/JP00/06375

33

amino which may have substituents" can be used.

Examples of "C₆₋₁₄ aryloxy-carbonyl" in "C₆₋₁₄ aryloxy-carbonyl which may have substituents" include phenyloxycarbonyl, 1-naphthyloxycarbonyl, 2-naphthyloxycarbonyl.

Examples of "C₇₋₁₉ aralkyloxy-carbonyl" in "C₇₋₁₉ aralkyloxy-carbonyl which may have substituents" include benzyloxycarbonyl, phenethyloxycarbonyl, diphenylmethyloxycarbonyl, triphenylmethyloxycarbonyl, 1-naphthylmethyloxycarbonyl, 2-naphthylmethyloxycarbonyl, 2,2-diphenylethyloxycarbonyl, 3-phenylpropyloxycarbonyl, 4-phenylbutyloxycarbonyl, 5-phenylpentyloxycarbonyl.

Examples of "5 to 6 membered hetero ring-carbonyl" in the above "5 to 6 membered hetero ring-carbonyl which may have substituents" include nicotinoyl, isonicotinoyl, 2-thenoyl, 3-thenoyl, 2-furoyl, 3-furoyl, morpholinocarbonyl, piperidinocarbonyl, pyrrolidin-1-ylcarbonyl.

Examples of the "5 to 6 membered hetero ring-carbamoyl" in the above "5 to 6 membered hetero ring-carbamoyl which may have substituents" include morpholinocarbamoyl, piperidinocarbamoyl, 2-pyridylcarbamoyl, 3-pyridylcarbamoyl, 4-pyridylcarbamoyl, 2-thienylcarbamoyl, 3-thienylcarbamoyl.

Examples of "C₆₋₁₄ arylsulfonyl" in the above "C₆₋₁₄ arylsulfonyl which may have substituents" include phenylsulfonyl, 1-naphthylsulfonyl, 2-naphthylsulfonyl.

Examples of "C₆₋₁₄ aryl-carbonyloxy" in the above "C₆₋₁₄ aryl-carbonyloxy which may have substituents" include benzoyloxy, 1-naphthoyloxy, 2-naphthoyloxy.

Examples of "C₆₋₁₄ aryl-carbamoyloxy" in the above "C₆₋₁₄ aryl-carbamoyloxy which may have substituents" include phenylcarbamoyloxy, naphthylcarbamoyloxy.

As the "substituents" in the above "C₆₋₁₄ aryloxy-

carbonyl which may have substituents", "C₇₋₁₉ aralkyloxy-carbonyl which may have substituents", "5 to 6 membered hetero ring-carbonyl which may have substituents", "5 to 6 membered hetero ring-carbamoyl which may have substituents", "C₆₋₁₄ arylsulfonyl which may have substituents", "C₆₋₁₄ aryl-carbonyloxy which may have substituents" and "C₆₋₁₄ aryl-carbamoyloxy which may have substituents", those exemplified as "substituents" in the above "C₇₋₁₉ aralkyl which may have substituents" can be mentioned. The number of the substituents is, for instance, 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituents can be the same or different.

Examples of "heterocyclic groups" in the "heterocyclic groups which may have substituents" for R³ or R^{3a} include a 5 to 14 membered (monocyclic, bicyclic or tricyclic) hetero ring containing 1 or 2 kinds of, 1 to 4 hetero atoms selected from nitrogen, sulfur and oxygen atom in addition to carbon atoms. Preferably, univalent groups formed by removing an optional one hydrogen atom from (i) an aromatic hetero ring, (ii) a 5 to 10 membered non-aromatic hetero ring, or (iii) a 7 to 10 membered hetero-bridge ring, can be mentioned.

Here, examples of the "aromatic hetero ring" include a 5 to 14 membered, preferably 5 to 10 membered, aromatic hetero ring containing one or more hetero atom (e.g. 1 to 4) selected from nitrogen, sulfur and oxygen atom in addition to carbon atoms.

Concrete examples include aromatic hetero rings such as thiophene, furan, pyrrole, imidazole, pyrazole, thiazole, isothiazole, oxazole, isoxazole, pyridine, pyrazine, pyrimidine, pyridazine, 1,2,4-oxadiazole, 1,3,4-oxadiazole, 1,2,4-thiadiazole, 1,3,4-thiadiazole, furazan, benzothiophene, benzofuran, benzimidazole, benzoxazole, benzothiazole, benzisothiazole,

naphtho[2,3-b]thiophene, phenoxathiin, indole, isoindole, 1H-indazole, purine, 4H-quinolidine, isoquinoline, quinoline, phthalazine, naphthylidine, quinoxaline, quinazoline, cinnoline, carbazole, β -carboline, phenanthridine, acridine, phenazinephenothiadine, phenoxazine, phthalimide, etc.; or a ring formed by condensing these rings (preferably monocyclic rings) with one to multiple (preferably 1 or 2) aromatic rings (e.g. benzene ring, etc.).

10 Examples of "5 to 10 membered non-aromatic hetero rings" include 2- or 3-pyrroline, pyrrolidine, 2- or 3-imidazoline, 2-oxazoline, oxazolidine, 2- or 3-pyrazoline, pyrazolidine, 2-thiazoline, piperidine, piperazine, hexamethylenimine, morpholine, thiomorpholine.

15 Examples of "7 to 10 membered hetero-bridge rings" include quinuclidine, 7-azabicyclo[2.2.1]heptane.

The "hetero cyclic groups" are preferably 5 to 10 membered (monocyclic or bicyclic) heterocyclic groups containing 1 or 2 kinds of, preferably 1 to 4, hetero atoms selected from nitrogen, sulfur and oxygen atom in addition to carbon atoms. Concretely examples include aromatic heterocyclic groups such as 2- or 3-thienyl; 2-, 3- or 4-pyridyl; 2- or 3-furyl; 2-, 4- or 5-thiazolyl; 2-, 4- or 5-oxazolyl; 1-, 3- or 4-pyrazolyl; 2-pyrazinyl; 2-, 4- or 5-pyrimidinyl; 1-, 2- or 3-pyrrolyl; 1-, 2- or 4-imidazolyl; 3- or 4-pyridazinyl; 3-isothiazolyl; 3-isoxazolyl; 1,2,4-oxadiazol-5-yl; 1,2,4-oxadiazol-3-yl; 2-, 3-, 4-, 5- or 8-quinolyl; 1-, 3-, 4-, 5-, 6-, 7- or 8-isoquinolyl; 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl; 1-, 2-, 4- or 5-isoindolyl; 1-, 5- or 6-phthalazinyl; 2-, 3- or 5-quinoxalinyl; 2-, 3-, 4-, 5- or 6-benzofuranyl; 2-, 3-, 4-, 5- or 6-benzothienyl; 2-, 4-, 5- or 6-benzothiazolyl; 1-, 2-, 4-, 5- or 6-benzimidazolyl; and non-aromatic heterocyclic groups such as 1-, 2- or 3-pyrrolidinyl; 1-, 2-, 4- or 5-imidazolidinyl; 2- or 4-imidazolinyl; 2-, 3- or 4-pyrazolidinyl; piperidino; 2-, 3- or 4-piperidyl; 1-

WO 01/21577

PCT/JP00/06375

36

or 2-piperazinyl; morpholino.

As the "substituents" in the "heterocyclic groups which may have substituents", those exemplified as "substituents" in the above "5 to 10 membered aromatic
5 heterocyclic groups which may have substituents" can be used. The number of substituents is, for instance, 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituents can be the same or different.

Examples of "C₁₋₆ alkyl" for R⁴ include methyl, ethyl,
10 propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, hexyl.

Examples of "nitrogen-containing hetero ring" in the "nitrogen-containing hetero ring which may have substituents" formed by R³ and R⁴ together with the adjacent
15 nitrogen atoms, include a 5 to 7 membered nitrogen-containing hetero ring which contains at least one nitrogen atom in addition to carbon atoms and may contain 1 to 3 hetero atoms selected from nitrogen, sulfur and oxygen atom. The "nitrogen-containing hetero rings" are preferably
20 piperidine, morpholine, thiomorpholine, piperazine, pyrrolidine, etc.

As the "substituents" in the "nitrogen-containing hetero ring which may have substituents", those exemplified as "substituents" in the above "5 to 10 membered aromatic
25 heterocyclic groups which may have substituents" can be used. The number of substituents is, for instance, 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituents can be the same or different.

The "acyl" is preferably formyl, carboxy, carbamoyl,
30 optionally halogenated C₁₋₆ alkyl-carbonyl (e.g. acetyl, etc.), C₁₋₆ alkoxy-carbonyl (e.g. methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, tert-butoxycarbonyl, etc.), C₆₋₁₄ aryl-carbonyl which may have substituents (e.g. benzoyl, 1-naphthoyl, 2-naphthoyl, etc.), C₆₋₁₄ aryloxy-carbonyl which may have substituents (e.g.
35 phenyloxycarbonyl, 1-naphthyloxycarbonyl, 2-

WO 01/21577

PCT/JP00/06375

37

naphthyloxycarbonyl, etc.), C_{7-19} aralkyloxy-carbonyl which may have substituents (e.g. benzyloxycarbonyl, phenethyloxycarbonyl, etc.), a 5 to 6 membered hetero ring-carbonyl which may have substituents (e.g. nicotinoyl, etc.), mono- C_{1-6} alkyl-carbamoyl (e.g. methylcarbamoyl, ethylcarbamoyl, etc.), di- C_{1-6} alkyl-carbamoyl (e.g. dimethylcarbamoyl, diethylcarbamoyl, ethylmethylcarbamoyl, etc.), C_{6-14} aryl-carbamoyl which may have substituents (e.g. phenylcarbamoyl, 4-methoxyphenylcarbamoyl, 3,4-dimethoxyphenylcarbamoyl, etc.), aromatic hetero ring-carbamoyl which may have substituents (e.g. 2-pyridinylcarbamoyl, 2-quinolinylcarbamoyl etc.), optionally halogenated C_{1-6} alkylsulfonyl (e.g. methylsulfonyl, etc.), C_{6-14} arylsulfonyl which may have substituents (e.g. phenylsulfonyl etc.), etc.

Here, as "optionally halogenated C_{1-6} alkyl-carbonyl" and "optionally halogenated C_{7-19} aralkylsulfonyl", those exemplified as "substituents" in the above " C_{7-19} aralkyl which may have substituents" can be used.

As " C_{6-14} aryl-carbonyl which may have substituents", "substituents" in the above "5 to 7 membered saturated cyclic amino which may have substituents" can be used.

As " C_{6-14} aryloxy-carbonyl which may have substituents", " C_{7-19} aralkyloxy-carbonyl which may have substituents", "5 to 6 membered hetero ring-carbonyl which may have substituents", "aromatic hetero ring-carbamoyl which may have substituents" and " C_{6-14} arylsulfonyl which may have substituents", those exemplified as "substituents" in the above "hydrocarbon groups which may have substituents" can be used.

As " C_{6-14} aryl-carbamoyl which may have substituents", those exemplified as "substituents" in the above "cyclic group which may have substituents" can be used.

35

Examples of the above "acylamino" include amino which

is substituted by 1 or 2 of the above "acyl". Preferably, acylamino of the formulae : $-NR^5-COR^6$, $-NR^5-COOR^{6a}$, $-NR^5-SO_2R^{6a}$, $-NR^5-CONR^{6a}R^{6b}$, $-PO(-OR^5)-OR^6$, or $-PO_2-R^6$ wherein R^5 is hydrogen atom or C_{1-6} alkyl; R^6 has the same meaning as the above R^3 ; R^{6a} has the same meaning as the above R^{3a} ; and R^{6b} has the same meaning as R^4], can be mentioned.

As " C_{1-6} alkyl" for R^5 , the same one as in " C_{1-6} alkyl" for the above R^4 can be mentioned.

The "acylamino" is preferably formylamino, optionally halogenated C_{1-6} alkyl-carboxamide (e.g. methylcarboxamide, trifluoromethylcarboxamide, isopropylcarboxamide, etc.), C_{6-14} aryl-carboxamide which may have substituents (e.g. phenylcarboxamide, 2-methoxyphenylcarboxamide, 4-methoxyphenylcarboxamide, etc.), N-(C_{6-14} aryl-carbonyl which may have substituents)-N- C_{1-6} alkylamino (e.g. N-4-methoxybenzoyl-N-methylamino, etc.), C_{7-19} aralkyl-carboxamide which may have substituents (e.g. benzylcarboxamide, etc.), aromatic hetero ring-carboxamide which may have substituents (e.g. benzothiophen-2-ylcarboxamide, etc.), optionally halogenated C_{1-6} alkoxy-carboxamide (e.g. methoxycarboxamide, ethoxycarboxamide, propoxycarboxamide, butoxycarboxamide, etc.), C_{6-14} arylamino-carbonylamino which may have substituents (e.g. phenylaminocarbonylamino, etc.), optionally halogenated C_{1-6} alkylsulfonylamino (e.g. methylsulfonylamino, trifluoromethylsulfonylamino, ethylsulfonylamino, etc.), C_{6-14} arylsulfonylamino which may have substituents (e.g. 4-methoxyphenylsulfonylamino, etc.).

Here, as "substituents" in " C_{6-14} aryl-carboxamide which may have substituents", "N-(C_{6-14} aryl-carbonyl which may have substituents)-N- C_{1-6} arylkylamino", " C_{7-19} aralkyl-carboxamide which may have substituents", "aromatic hetero ring-carboxamide which may have substituents", " C_{6-14} arylamino-carbonylamino which may

WO 01/21577

PCT/JP00/06375

39

have substituents" and "C₆₋₁₄ arylsulfonylamino which may have substituents", those exemplified as "substituents" in the above "C₇₋₁₉ aralkyl which may have substituents" can be mentioned. The number of substituents is, for instance,
5 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituents can be the same or different.

Examples of the above "acyloxy" include oxy substituted by one of the above "acyl". Preferably,
10 acyloxy of the formulae : -O-COR⁷, -O-COOR⁷, -O-CONHR⁷, -PO(OH)-OR⁷ or -PO₂-R⁷ wherein R⁷ has the same meaning as the above R³, can be mentioned.

The "acyloxy" is preferably optionally halogenated C₁₋₆ alkyl-carbonyloxy (e.g. acetoxy, propanoyloxy, etc.),
15 C₆₋₁₄ aryl-carbonyloxy which may have substituents (e.g. benzoyloxy, 4-methoxybenzoyloxy, etc.), optionally halogenated C₁₋₆ alkoxy-carbonyloxy (e.g. methoxycarbonyloxy, trifluoromethoxycarbonyloxy, ethoxycarbonyloxy, propoxycarbonyloxy,
20 butoxycarbonyloxy, etc.), mono-C₁₋₆ alkyl-carbamoyloxy (e.g. methylcarbamoyloxy, ethylcarbamoyloxy, etc.), di-C₁₋₆ alkyl-carbamoyloxy (e.g. dimethylcarbamoyloxy, diethylcarbamoyloxy, etc.), C₆₋₁₄ aryl-carbamoyloxy which may have substituents (e.g. phenylcarbamoyloxy,
25 naphthylcarbamoyloxy, etc.), nicotinyloxy, etc.

As "substituents" in "C₆₋₁₄ aryl-carbonyloxy which may have substituents" and "C₆₋₁₄ aryl-carbamoyloxy which may have substituents", those exemplified as "substituents" in the above "C₇₋₁₉ aralkyl which may have substituents" can
30 be mentioned. The number of substituents is, for instance, 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituents can be the same or different.

Examples of the "5 to 7 membered non-aromatic
35 heterocyclic groups which may have substituents", which is "substituents" in "cyclic group which may have

substituents" for Ar¹, include 4,5-dihydro-1,3-oxazol-2-yl, 4,5-dihydro-1,3-thiazol-2-yl, 4,5-dihydro-1H-2-imidazolyl. As "substituents" in the "5 to 7 membered non-aromatic heterocyclic groups which may have substituents", those exemplified as "substituents" in the above "5 to 7 membered saturated cyclic amino which may have substituents" can be used.

As "acyl", "acyloxy" and "acylamino", which are "substituents" in the "cyclic group which may have substituents" for Ar¹, those exemplified as "substituents" in the above "5 to 10 membered aromatic heterocyclic groups which may have substituents" can be used.

Regarding "aromatic hetero ring-C₁₋₆ alkoxy" which is "substituents" in the "cyclic group which may have substituents" for Ar¹, as "aromatic hetero ring", those exemplified as the above R³ can be used. Examples of "C₁₋₆ alkoxy" include methoxy, ethoxy, propoxy, butoxy, pentyloxy.

"Substituents" in the "cyclic group which may have substituents" for Ar¹ are preferably halogen atom (preferably fluorine, chlorine and bromine, etc.); nitro; C₁₋₃ alkylenedioxy (preferably methylenedioxy, etc.); optionally halogenated C₁₋₆ alkyl (preferably, methyl, ethyl, propyl, trifluoromethyl, etc.); hydroxy-C₁₋₆ alkyl (preferably hydroxymethyl, etc.); optionally halogenated C₃₋₆ cycloalkyl (preferably cyclohexyl, etc.); optionally halogenated C₁₋₆ alkoxy (preferably methoxy, ethoxy, etc.); optionally halogenated C₁₋₆ alkylthio (preferably methylthio, etc.); hydroxy; C₇₋₁₉ aralkyloxy which may have substituents (preferably, 1 to 3 substituents selected from halogen atom, optionally halogenated C₁₋₆ alkyl, optionally halogenated C₁₋₆ alkoxy, optionally halogenated C₁₋₆ alkylthio, etc.) (preferably benzyloxy, 4-methoxybenzyloxy, 3-methoxybenzyloxy, 4-fluorobenzyloxy, 4-methylthiobenzyloxy, 4-ethylbenzyloxy, etc.); C₆₋₁₄

WO 01/21577

PCT/JP00/06375

41

aryloxy which may have substituents (preferably, 1 to 3 optionally halogenated C₁₋₆ alkoxy, etc.) (preferably phenyloxy, 4-methoxyphenyloxy, etc.); amino; mono-C₁₋₆ alkylamino (preferably methylamino, etc.); di-C₁₋₆ alkylamino (preferably dimethylamino, etc.); 5 to 7 membered saturated cyclic amino which may have substituents (preferably 1 to 3 oxo) and may be condensed with a benzene ring (preferably 1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl, etc.); 5 to 7 membered non-aromatic heterocyclic groups which may have substituents (preferably 4,5-dihydro-1,3-oxazol-2-yl, etc.); formyl; carboxy; C₆₋₁₄ aryl-carbonyl which may have substituents (preferably benzoyl, etc.); C₆₋₁₄ aryl-carbamoyl which may have substituents (preferably, 1 to 3 optionally halogenated C₁₋₆ alkoxy, etc.) (preferably, phenylcarbamoyl, 4-methoxyphenylcarbamoyl, 3,4-dimethoxyphenylcarbamoyl, etc.); aromatic hetero ring-carbamoyl which may have substituents (preferably 2-pyridinylcarbamoyl, 2-quinolinylcarbamoyl, etc.); C₁₋₆ alkoxy-carbonyl (preferably methoxycarbonyl, ethoxycarbonyl, etc.); optionally halogenated C₁₋₆ alkyl-carboxamide (preferably methylcarboxamide, trifluoromethylcarboxamide, isopropylcarboxamide, etc.); C₆₋₁₄ aryl-carboxamide which may have substituents (preferably, 1 to 3 optionally halogenated C₁₋₆ alkoxy, etc.) (preferably phenylcarboxamide, 2-methoxyphenylcarboxamide, 4-methoxyphenylcarboxamide, etc.); C₇₋₁₉ aralkyl-carboxamide which may have substituents (preferably benzylcarboxamide, etc.); aromatic hetero ring-carboxamide which may have substituents (preferably benzothiophen-2-ylcarboxamide, etc.); N-(C₆₋₁₄ aryl-carbonyl which may have substituents (preferably, 1 to 3 optionally halogenated C₁₋₆ alkoxy, etc.))-N-C₁₋₆ alkylamino (preferably N-4-methoxybenzoyl-N-methylamino, etc.); C₆₋₁₄ arylamino-carbonylamino which may have substituents (preferably phenylaminocarbonylamino, etc.); C₆₋₁₄ arylsulfonylamino which may have substituents (preferably,

1 to 3 optionally halogenated C₁₋₆ alkoxy, etc.) (preferably 4-methoxyphenylsulfonylamino, etc.); C₆₋₁₄ aryl-carbonyloxy which may have substituents (preferably, 1 to 3 optionally halogenated C₁₋₆ alkoxy, etc.) (preferably 4-methoxybenzoyloxy, etc.); oxo; carboxy-C₁₋₆ alkyl (preferably carboxyethyl, etc.); C₁₋₆ alkoxy-carbonyl-C₁₋₆ alkyl (preferably methoxycarbonylmethyl, etc.); C₇₋₁₉ aralkyl which may have substituents (preferably 1 to 3 halogen atom) (preferably benzyl, 2,4-dichlorobenzyl, etc.); aromatic hetero ring-C₁₋₆ alkoxy (preferably 2-guanolylmethoxy, etc.); cyano, etc.

When "cyclic group" in "cyclic group which may have substituents" for Ar¹ is a non-aromatic cyclic hydrocarbon group or a non-aromatic heterocyclic group, C₆₋₁₄ aryl which may have substituents (preferably, 1 to 3 substituents selected from halogen atom, C₁₋₃ alkylenedioxy, optionally halogenated C₁₋₆ alkyl, optionally halogenated C₁₋₆ alkoxy, etc.) (preferably phenyl, 4-fluorophenyl, 1,3-benzodioxol-5-yl, 4-chlorophenyl, 4-methylphenyl, 4-methoxyphenyl), hydroxy, C₇₋₁₉ aralkyloxy-carbonyl (preferably benzyloxycarbonyl), C₇₋₁₉ aralkyl (preferably benzyl), etc., can be used as a preferable substituent.

Ar¹ is preferably phenyl, biphenyl (preferably 4-biphenyl, 2-biphenyl), phenyl-pyridyl (preferably 6-phenyl-3-pyridyl, 5-phenyl-2-pyridyl), phenyl-furyl (preferably 5-phenyl-2-furyl), phenyl-isoxazolyl (preferably 3-phenyl-isoxazol-5-yl), diphenyl-oxazolyl (preferably 2,4-diphenyl-1,3-oxazol-5-yl), pyridyl-phenyl (preferably 4-(4-pyridyl)phenyl, 4-(3-pyridyl)phenyl), phenyl-pyrimidinyl (preferably 2-phenyl-5-pyrimidinyl), benzofuranyl-phenyl (preferably 4-(2-benzofuranyl)phenyl), furyl-phenyl (preferably 4-(2-furyl)phenyl), terphenyl (preferably 4,4'-terphenyl), thienyl-phenyl (preferably 4-(2-thienyl)phenyl), indolyl (preferably 2-indolyl, 3-indolyl), naphthyl-oxadiazolyl

WO 01/21577

PCT/JP00/06375

43

(preferably 3-(2-naphthyl)-1,2,4-oxadiazol-5-yl),
benzofuranyl-oxadiazole (preferably 3-(2-benzofuranyl)-
1,2,4-oxadiazol-5-yl), benzothienyl (preferably 2-
benzothienyl), benzofuranyl (preferably 2-benzofuranyl),
5 fluorenyl (preferably 2-fluorenyl), pyridyl-pyrrolyl
(preferably 3-(4-pyridyl)pyrrolyl), thioxanthenyl;
each of which may have 1 to 3 (preferably 1 or 2) substituents
selected from the group consisting of halogen atom
(preferably fluorine, chlorine, bromine, etc.); nitro; C₁₋₃
10 alkylenedioxy (preferably methylenedioxy, etc.);
optionally halogenated C₁₋₆ alkyl (preferably methyl,
ethyl, propyl, trifluoromethyl, etc.); hydroxy-C₁₋₆ alkyl
(preferably hydroxymethyl, etc.); optionally halogenated
C₃₋₆ cycloalkyl (preferably cyclohexyl, etc.); optionally
15 halogenated C₁₋₆ alkoxy (preferably methoxy, ethoxy, etc.);
optionally halogenated C₁₋₆ alkythio (preferably
methylthio, etc.); hydroxy; C₇₋₁₉ aralkyloxy which may have
substituents (preferably, 1 to 3 substituents selected from
halogen atom, optionally halogenated C₁₋₆ alkyl, optionally
20 halogenated C₁₋₆ alkoxy, optionally halogenated C₁₋₆
alkylthio, etc.) (preferably benzyloxy, 4-
methoxybenzyloxy, 3-methoxybenzyloxy, 4-fluorobenzyloxy,
4-methylthiobenzyloxy, 4-ethylbenzyloxy, etc.); C₆₋₁₄
aryloxy which may have substituents (preferably, 1 to 3
25 optionally halogenated C₁₋₆ alkoxy, etc.) (preferably
phenyloxy, 4-methoxyphenyloxy, etc.); amino; mono-C₁₋₆
alkylamino (preferably methylamino, etc.); di-C₁₋₆
alkylamino (preferably dimethylamino, etc.); 5 to 7
membered saturated cyclic amino which may have substituents
30 (preferably 1 to 3 oxo) and may be condensed with a benzene
ring (preferably 1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl,
etc.); 5 to 7 membered non-aromatic heterocyclic groups
which may have substituents (preferably 4,5-dihydro-
1,3-oxazol-2-yl, etc.); formyl; carboxy; C₆₋₁₄ aryl-
35 carbonyl which may have substituents (preferably benzoyl,
etc.); C₆₋₁₄ aryl-carbamoyl which may have substituents

WO 01/21577

PCT/JP00/06375

44

(preferably, 1 to 3 optionally halogenated C₁₋₆ alkoxy, etc.)
(preferably phenylcarbamoyl, 4-methoxyphenylcarbamoyl,
3,4-dimethoxyphenylcarbamoyl, etc.); aromatic hetero
ring-carbamoyl which may have substituents (e.g. 2-
5 pridinylicarbamoyl, 2-quinolinylcarbamoyl, etc.); C₁₋₆
alkoxy-carbonyl (preferably methoxycarbonyl,
ethoxycarbonyl, etc.); optionally halogenated C₁₋₆
alkyl-carboxamide (preferably, methylcarboxamide,
trifluoromethylcarboxamide, isopropylcarboxamide, etc.);
10 C₆₋₁₄ aryl-carboxamide which may have substituents
(preferably, 1 to 3 optionally halogenated C₁₋₆ alkoxy, etc.)
(preferably phenylcarboxamide, 2-
methoxyphenylcarboxamide, 4-methoxyphenylcarboxamide,
etc.); C₇₋₁₉ aralkyl-carboxamide which may have substituents
15 (preferably benzylcarboxamide, etc.); aromatic hetero
ring-carboxamide which may have substituents (preferably
benzothiophen-2-ylcarboxamide, etc.); N-(C₆₋₁₄ aryl-
carbonyl which may have substituents (preferably, 1 to 3
optionally halogenated C₁₋₆ alkoxy, etc.))-N-C₁₋₆ alkylamino
20 (preferably N-4-methoxybenzoyl-N-methylamino, etc.); C₆₋₁₄
arylamino-carbonylamino which may have substituents
(preferably phenylaminocarbonylamino, etc.); C₆₋₁₄
arylsulfonylamino which may have substituents (preferably,
1 to 3 optionally halogenated C₁₋₆ alkoxy, etc.) (preferably
25 4-methoxyphenylsulfonylamino, etc.); C₆₋₁₄ aryl-
carbonyloxy which may have substituents (preferably, 1 to
3 optionally halogenated C₁₋₆ alkoxy, etc.) (preferably
4-methoxybenzoyloxy, etc.); oxo; carboxy-C₁₋₆ alkyl
(preferably carboxyethyl, etc.); C₁₋₆ alkoxy-carbonyl-C₁₋₆
30 alkyl (preferably methoxycarbonylmethyl, etc.); C₇₋₁₉
aralkyl which may have substituents (preferably 1 to 3
halogen atom) (preferably benzyl, 2,4-dichlorobenzyl,
etc.); aromatic hetero ring-C₁₋₆ alkoxy (preferably 2-
quinolylmethoxy, etc.); and cyano.

35 Further, preferable examples of Ar¹ include
piperidiny (preferably piperidino), piperaziny,

pyrrolidinyl, dihydropyridyl, tetrahydropyridyl; each of which may have 1 or 2 substituents selected from the group consisting of oxo, C₆₋₁₄ aryl which may have substituents (preferably, 1 to 3 substituents selected from halogen atom, C₁₋₃ alkylenedioxy, optionally halogenated C₁₋₆ alkyl, optionally halogenated C₁₋₆ alkoxy, etc.) (preferably phenyl, 4-fluorophenyl, 1,3-benzodioxol-5-yl, 4-chlorophenyl, 4-methylphenyl, 4-methoxyphenyl), hydroxy, C₇₋₁₉ aralkyloxy-carbonyl (preferably benzyloxycarbonyl) and C₇₋₁₉ aralkyl (preferably benzyl).

Ar¹ is more preferably, phenyl, biphenyl (preferably 4-biphenyl) or phenyl-pyridyl (preferably 6-phenyl-3-pyridyl, 5-phenyl-2-pyridyl); each of which may have 1 or 2 substituents selected from the group consisting of halogen atom (preferably fluorine, chlorine, bromine, etc.); optionally halogenated C₁₋₆ alkyl (preferably methyl, ethyl, propyl, trifluoromethyl, etc.); optionally halogenated C₁₋₆ alkoxy (preferably methoxy, ethoxy, etc.); C₇₋₁₉ aralkyloxy which may have substituents (preferably, 1 to 3 substituents selected from halogen atom, optionally halogenated C₁₋₆ alkyl, optionally halogenated C₁₋₆ alkoxy, optionally halogenated C₁₋₆ alkylthio, etc.) (preferably benzyloxy, 4-methoxybenzyloxy, etc.); C₆₋₁₄ aryloxy which may have substituents (preferably, 1 to 3 optionally halogenated C₁₋₆ alkoxy, etc.) (preferably phenoxy, etc.); C₆₋₁₄ aryl-carbonyl which may have substituents (preferably, 1 to 3 optionally halogenated C₁₋₆ alkoxy, etc.) (preferably benzoyl, etc.); C₆₋₁₄ aryl-carbamoyl which may have substituents (preferably, 1 to 3 optionally halogenated C₁₋₆ alkoxy, etc.) (preferably phenylcarbamoyl, 4-methoxyphenylcarbamoyl, 3,4-dimethoxyphenylcarbamoyl, etc.); aromatic hetero ring-carbamoyl which may have substituents (e.g. 2-pyridinylcarbamoyl, 2-quinolinylcarbamoyl, etc.); C₆₋₁₄ aryl-carboxamide which may have substituents (preferably, 1 to 3 optionally halogenated C₁₋₆ alkoxy, etc.) (preferably

phenylcarboxamide, 2-methoxyphenylcarboxamide, 4-methoxyphenylcarboxamide, etc.); C₇₋₁₉ aralkyl-carboxamide which may have substituents (preferably benzylcarboxamide, etc.); aromatic hetero ring-carboxamide (preferably benzothiophen-2-ylcarboxamide, etc.); N-(C₆₋₁₄ aryl-carbonyl which may have substituents (preferably, 1 to 3 optionally halogenated C₁₋₆ alkoxy, etc.))-N-C₁₋₆ alkylamino (preferably N-4-methoxybenzoyl-N-methylamino, etc.); C₆₋₁₄ arylamino-carbonylamino which may have substituents (preferably phenylaminocarbonylamino, etc.); C₆₋₁₄ arylsulfonylamino which may have substituents (preferably, 1 to 3 optionally halogenated C₁₋₆ alkoxy, etc.) (preferably 4-methoxyphenylsulfonylamino, etc.); and C₆₋₁₄ arylcarbonyloxy which may have substituents (preferably, 1 to 3 optionally halogenated C₁₋₆ alkoxy, etc.) (preferably 4-methoxybenzoyloxy, etc.).

Further, preferable examples of Ar¹ include piperidino, piperazinyl or pyrrolidinyl; each of which may have 1 or 2 substituents selected from the group consisting of oxo and C₆₋₁₄ aryl (preferably phenyl) which may have substituents [preferably halogen atom (preferably fluorine, chlorine, bromine, etc.), optionally halogenated C₁₋₆ alkyl (preferably methyl, ethyl, propyl, trifluoromethyl, etc.) or optionally halogenated C₁₋₆ alkoxy (preferably methoxy, ethoxy, etc.)].

The "spacer having a main chain of 1 to 6 atoms" means a space in which 1 to 6 atoms are linked. Here, the "number of atoms in the main chain" is counted so that the number of atoms in the main chain is minimum. For instance, the number of atoms of 1,2-cyclopentylene is counted as 2, and the number of atoms of 1,3-cyclopentylene is counted as 3.

Examples of the "spacer having a main chain of 1 to 6 atoms" include a bivalent group consisting of 1 to 3 species selected from -O-, -S-, -CO-, -SO-, -SO₂-, -NR⁸- (R⁸ is hydrogen atom, optionally halogenated C₁₋₆ alkyl,

WO 01/21577

PCT/JP00/06375

47

optionally halogenated C₁₋₆ alkyl-carbonyl, optionally halogenated C₁₋₆ alkylsulfonyl), bivalent C₁₋₆ non-cyclic hydrocarbon groups which may have substituents, and bivalent C₅₋₈ monocyclic non-aromatic hydrocarbon groups.

5 Here, as "optionally halogenated C₁₋₆ alkyl", those exemplified as "substituents" in the above "cyclic group which may have substituents" can be used.

As "optionally halogenated C₁₋₆ alkyl-carbonyl" and "optionally halogenated C₁₋₆ alkylsulfonyl", those
10 exemplified as "substituents" in the above "C₇₋₁₉ aralkyl which may have substituents" can be used.

Examples of "bivalent C₁₋₆ non-cyclic hydrocarbon groups" in the "bivalent C₁₋₆ non-cyclic hydrocarbon groups which may have substituents" include
15

(1) C₁₋₆ alkylene (e.g. -CH₂-, -(CH₂)₂-, -(CH₂)₃-, -(CH₂)₄-, -(CH₂)₅-, -(CH₂)₆-, -CH(CH₃)-, -C(CH₃)₂-, -(CH(CH₃))₂-, -(CH₂)₂C(CH₃)₂-, -(CH₂)₃C(CH₃)₂-, etc.);

(2) C₂₋₆ alkenylene (e.g. -CH=CH-, -CH₂-CH=CH-, -C(CH₃)₂-CH=CH-, -CH₂-CH=CH-CH₂-, -CH₂-CH₂-CH=CH-, -CH=CH-CH=CH-, -CH=CH-CH₂-CH₂-CH₂-, etc.);
20

(3) C₂₋₆ alkynylene (e.g. -C≡C-, -CH₂-C≡C-, -CH₂-C≡C-CH₂-CH₂-, etc.)

each of which may have 1 to 5, preferably 1 to 3, halogen atoms (e.g. fluorine, chlorine, bromine, iodine, etc.).
25

The "bivalent C₁₋₆ non-cyclic hydrocarbon groups" may have 1 to 5, preferably 1 to 3 substituents at a substitutable position. Examples of such substituents include halogen atom (e.g. fluorine, chlorine, bromine, iodine, etc.), hydroxy, C₁₋₆ alkyl-carbonyloxy (e.g.,
30 acetoxo, etc.).

As the "bivalent C₅₋₈ monocyclic non-aromatic hydrocarbon groups", for instance, bivalent groups formed
35 by removing an optional two hydrogen atoms from C₅₋₈ cycloalkane or C₅₋₈ cycloalkene, can be mentioned. Concrete

WO 01/21577

PCT/JP00/06375

48

examples include 1,2-cyclopentylene; 1,3-cyclopentylene;
1,2-cyclohexylene; 1,3-cyclohexylene; 1,4-cyclohexylene;
1,2-cycloheptylene; 1,3-cycloheptylene; 1,4-
cycloheptylene; 3-cyclohexen-1,4-ylene; 3-cyclohexen-
5 1,2-ylene; 2,5-cyclohexadien-1,4-ylene. Especially, C₅₋₈
cycloalkylene is preferable.

The "spacer having a main chain of 1 to 6 atoms" is
preferably a bivalent group consisting of 1 to 3 species
selected from -O-, -S-, -CO-, -SO-, -SO₂-, -NR⁸- (R⁸ has
10 the same meaning as defined above) and optionally
halogenated bivalent C₁₋₆ non-cyclic hydrocarbon groups.

Preferred examples of the "spacer having a main chain
of 1 to 6 atoms" include

(1) C₁₋₆ alkylene (e.g. -CH₂-, -(CH₂)₂-, -(CH₂)₃-, -
15 (CH₂)₄-, -(CH₂)₅-, -(CH₂)₆-, -CHCH₃-, -C(CH₃)₂-, -CH(CF₃)-,
-(CH(CH₃))₂-, -(CF₂)₂-, -(CH₂)₂C(CH₃)₂-, -(CH₂)₃C(CH₃)₂-,
etc.);

(2) C₂₋₆ alkenylene (e.g. -CH=CH-, -CH₂-CH=CH-, -
CH₂-CF=CH-, -C(CH₃)₂-CH=CH-, -CH₂-CH=CH-CH₂-, -CH₂-CH₂-
20 CH=CH-, -CH=CH-CH=CH-, -CH=CH-CH₂-CH₂-CH₂-, etc.);

(3) C₂₋₆ alkynylene (e.g. -C≡C-, -CH₂-C≡C-, -CH₂-C
≡C-CH₂-, etc.);

(4) -(CH₂)_{w1}O(CH₂)_{w2}-, -(CH₂)_{w1}S(CH₂)_{w2}-,
-(CH₂)_{w1}CO(CH₂)_{w2}-, -(CH₂)_{w1}SO(CH₂)_{w2}-,
25 -(CH₂)_{w1}SO₂(CH₂)_{w2}-, -(CH₂)_{w1}NR⁸(CH₂)_{w2}-;

(5) -(CH₂)_{w3}CONR⁸(CH₂)_{w4}-, -(CH₂)_{w3}NR⁸CO(CH₂)_{w4}-,
-(CH₂)_{w3}SO₂NR⁸(CH₂)_{w4}-, -(CH₂)_{w3}NR⁸SO₂(CH₂)_{w4}-,
-(CH₂)_{w3}COO(CH₂)_{w4}-;

(6) -(CH₂)_{w5}NR⁸CONR⁸(CH₂)_{w6}-;

30 (7) -(CH₂)_{w7}CONR⁸-(CH₂)_{w8}-CONR^{8b}-(CH₂)_{w9}-;
-CH=CH-CONR⁸-; -CH=CH-SO₂NR⁸-;

wherein R⁸ has the same meaning as defined above; R^{8b} has
the same meaning as R⁸; w1 and w2 is an integer of 0 to
5, and w1 + w2 is 0 to 5; w3 and w4 is an integer of 0 to
35 4, and w3 + w4 is 0 to 4; w5 and w6 is an integer of 0 to
3, and w5 + w6 is 0 to 3; w7, w8 and w9 is an integer of

WO 01/21577

PCT/JP00/06375

49

0 to 2, and $w_7 + w_8 + w_9$ is 0 to 2.

5 The "spacer having a main chain of 1 to 6 atoms" for X, is preferably $-(CH_2)_{w_1}O(CH_2)_{w_2}-$ (symbols have the same meaning as defined above), $-CONR^{8c}-$, $-NR^{8c}CO-$, $-CH=CH-CONR^{8c}-$, $-SO_2NR^{8c}-$ (R^8 is hydrogen atom or C_{1-6} alkyl); more preferably $-CONR^{8c}-$, $-NR^{8c}CO-$, $-CH=CH-CONR^{8c}-$, $-SO_2NR^{8c}-$ (R^8 has the same meaning as defined above); especially preferably $-CONH-$, $-NHCO-$, etc.

10 The "spacer having a main chain of 1 to 6 atoms" for Y, is preferably optionally halogenated bivalent C_{1-6} non-cyclic hydrocarbon groups, $-(CH_2)_{w_3}CONH(CH_2)_{w_4}-$, $-(CH_2)_{w_3}COO(CH_2)_{w_4}-$ (symbols have the same meaning as defined above); more preferably C_{1-3} alkylene (e.g. $-CH_2-$, $-(CH_2)_2-$,
15 $-(CH_2)_3-$, etc.), $-(CH_2)_{w_3}CONH(CH_2)_{w_4}-$, $-(CH_2)_{w_3}COO(CH_2)_{w_4}-$ (symbols have the same meaning as defined above); especially preferably C_{1-3} alkylene (e.g. $-CH_2-$, $-(CH_2)_2-$, $-(CH_2)_3-$, etc.), etc.

20 As "substituents" and "monocyclic aromatic rings" in "monocyclic aromatic rings which may be condensed with 4 to 8 membered non-aromatic rings, and may have further substituents" for Ar, those exemplified as "substituents" and "cyclic group" in the "cyclic group which may have
25 substituents" for the above Ar¹ can be used. The number of substituents is, for instance, 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituents can be the same or different.

30 The substituents are preferably formyl, optionally halogenated C_{1-6} alkyl-carbonyl, optionally halogenated C_{1-6} alkylsulfonyl, etc.

35 Here, as "optionally halogenated C_{1-6} alkyl-carbonyl" and "optionally halogenated C_{1-6} alkylsulfonyl", those exemplified as "substituents" in " C_{7-19} aralkyl which may have substituents" can be used respectively.

WO 01/21577

PCT/JP00/06375

50


Examples of "4 to 8 membered non-aromatic rings" in the "monocyclic aromatic rings which may be condensed with 4 to 8 membered non-aromatic rings, and may have further substituents" include C₄₋₈ monocyclic non-aromatic hydrocarbon rings, 4 to 8 membered monocyclic non-aromatic hetero rings.

Examples of the "C₄₋₈ monocyclic non-aromatic hydrocarbon rings" include C₄₋₈ cycloalkane and C₄₋₈ cycloalkene. Concrete examples include cyclobutane, cyclopentane, cyclohexane, cycloheptane, cyclooctane, cyclopentene, cyclohexene, cycloheptene. Especially, cyclopentane, cyclohexane, cyclobutane, etc. are preferable.

Examples of the "4 to 8 membered monocyclic non-aromatic hetero rings" include azetidine, pyrrolidine, pyrroline, pyrazolidine, 2- or 3-pyrazoline, imidazoline, piperidine, piperazine, azepine, azokane, oxane, oxine, oxepane, oxazolidine, 2-oxazoline, thiazolidine, 2-thioazoline, morpholine, thiomorpholine.

The above "4 to 8 membered non-aromatic rings" may have 1 to 3 substituents at a substitutable position. Examples of such substituents include optionally halogenated C₁₋₆ alkyl (e.g., methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, hexyl, etc.), cyano, hydroxy.


Regarding Ar, concrete examples of "monocyclic aromatic rings which may be condensed with 4 to 8 membered non-aromatic rings, and may have further substituents" include




WO 01/21577

PCT/JP00/06375

51




Ar is preferably benzene, pyridine, or rings of the
5 formulae :



wherein ----- is a single bond or double bond; each of m
and n is an integer of 1 to 4.

10 Ar is more preferably benzene, pyridine, rings of the
formulae :



15 As the "hydrocarbon groups which may have
substituents" for R¹ and R², those exemplified as the above R³

WO 01/21577

PCT/JP00/06375

52

can be used.

The "hydrocarbon groups which may have substituents" are preferably "C₁₋₆ alkyl which may have substituents".

Here, examples of "C₁₋₆ alkyl" in the "C₁₋₆ alkyl which
5 may have substituents" include methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, hexyl. Especially, methyl, ethyl, propyl, etc. are preferable.

Examples of "substituents" in the "C₁₋₆ alkyl which may
10 have substituents" include halogen atom (e.g. fluorine, chlorine, bromine, iodine, etc.), C₁₋₃ alkylenedioxy (e.g. methylenedioxy, ethylenedioxy etc.), nitro, cyano, optionally halogenated C₃₋₆ cycloalkyl, optionally halogenated C₁₋₆ alkoxy, optionally halogenated C₁₋₆
15 alkylthio, hydroxy, amino, mono-C₁₋₆ alkylamino (e.g. methylamino, ethylamino, propylamino, isopropylamino, butylamino, etc.), di-C₁₋₆ alkylamino (e.g. dimethylamino, diethylamino, dipropylamino, dibutylamino, ethylmethylamino, etc.), formyl, carboxy, carbamoyl,
20 thiocarbamoyl, optionally halogenated C₁₋₆ alkyl-carbonyl, optionally halogenated C₁₋₆ alkoxy-carbonyl (e.g. methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, tert-butoxycarbonyl, etc.), mono-C₁₋₆ alkyl-carbamoyl (e.g. methylcarbamoyl, ethylcarbamoyl, etc.), di-C₁₋₆ alkyl-
25 carbamoyl (e.g. dimethylcarbamoyl, diethylcarbamoyl, ethylmethylcarbamoyl etc.), optionally halogenated C₁₋₆ alkylsulfonyl, formylamino, optionally halogenated C₁₋₆ alkyl-carboxamide, C₁₋₆ alkoxy-carboxamide (e.g. methoxycarboxamide, ethoxycarboxamide,
30 propoxycarboxamide, butoxycarboxamide, etc.), C₁₋₆ alkylsulfonylamino (e.g. methylsulfonylamino, ethylsulfonylamino, etc.), C₁₋₆ alkyl-carbonyloxy (e.g. acetoxyl, propanoyloxy, etc.), C₁₋₆ alkoxy-carbonyloxy (e.g. methoxycarbonyloxy, ethoxycarbonyloxy,
35 propoxycarbonyloxy, butoxycarbonyloxy, etc.), mono-C₁₋₆ alkyl-carbamoyloxy (e.g. methylcarbamoyloxy,

ethylcarbamoyloxy, etc.), di-C₁₋₆ alkyl-carbamoyloxy (e.g. dimethylcarbamoyloxy, diethylcarbamoyloxy, etc.), and aromatic groups which may have substituents. The number of substituents is, for instance, 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituents can be the same or different.

Here, as "optionally halogenated C₃₋₆ cycloalkyl," "optionally halogenated C₁₋₆ alkoxy" and "optionally halogenated C₁₋₆ alkylthio", those exemplified as "substituents" in the above "cyclic group which may have substituents" can be used.

As "optionally halogenated C₁₋₆ alkyl-carbonyl," "optionally halogenated C₁₋₆ alkylsulfonyl" and "optionally halogenated C₁₋₆ alkyl-carboxamide", those exemplified as "substituents" in the above "C₇₋₁₉ aralkyl which may have substituents" can be used.

As "substituents" and "aromatic groups" in the "aromatic groups which may have substituents", those exemplified as "substituents" and "aromatic groups" in the "cyclic group which may have substituents" for the above Ar¹ can be used. The number of substituents is, for instance, 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituents can be the same or different.

Examples of "nitrogen-containing hetero rings" in the "nitrogen-containing hetero rings which may have substituents" formed by R¹ and R² together with the adjacent nitrogen atom, include 3 to 8 membered nitrogen-containing hetero rings which contain at least one nitrogen atom in addition to carbon atoms, and which may further contain 1 to 3 hetero atoms selected from nitrogen, sulfur and oxygen atom. Concrete examples include aziridine, azetidine, morpholine, thiomorpholine, piperidine, piperazine, pyrrolidine, hexamethyleneimine, heptamethyleneimine, hexahydropyrimidine, 1,4-diazepan, 4,5-dihydro-imidazole, and their unsaturated cyclic amines (e.g.

WO 01/21577

PCT/JP00/06375

54

1,2,5,6-tetrahydropyridine, etc.) can be mentioned.
Especially, morpholine, piperidine, piperazine,
pyrrolidine.


As "substituents" in the "nitrogen-containing hetero
rings which may have substituents", for instance, those
exemplified as "substituents" in the above "5 to 7 membered
saturated cyclic amino which may have substituents" can be
used. The number of substituents is, for instance, 1 to
5, preferably 1 to 3. When the number of substituents is
2 or more, each substituents can be the same or different.

R^1 and R^2 are preferably C_{1-6} alkyl, more preferably
methyl, ethyl, propyl, etc.


Also, it is preferable that R^1 and R^2 , together with
the adjacent nitrogen atom, form piperidino,
pyrrolidin-1-yl, piperazin-1-yl etc.

And, it is preferable that at least one of R^1 and R^2
is C_{1-6} alkyl which may have substituents. It is especially
preferable that both R^1 and R^2 is C_{1-6} alkyls which may have
substituents.

R^2 can form a spiro ring together with Ar. For
instance, Ar is a ring of the formula :



wherein n is an integer of 1 to 4; and Y is methylene; R^2
can form a spiro ring together with Ar. Examples of the
spiro ring include




wherein k (ring Ar and N are connected by $-(CH_2)_k-$) is an
integer of 1 to 4; and other symbols have the same meaning
as defined above.

Chemical structures of 15 compounds (1-15) used in the study. The structures are arranged in a grid. 1-3 are in the first row, 4-6 in the second, 7-9 in the third, 10-12 in the fourth, and 13-15 in the fifth. The structures include various combinations of naphthalene, tetralin, and indole systems, with different nitrogen-containing side chains such as dimethylamino, pyrrolidino, piperidino, morpholino, and N-benzyl groups.


WO 01/21577

PCT/JP00/06375

56



Among the compounds of the formula (I), a compound wherein Ar is a ring of the formula :




10 wherein ----- is a single bond or double bond, n is an integer of 1 to 4, and each ring may have substituents;

X is -CONR^{8c}-, -NR^{8c}CO-, -CH=CH-CONR^{8c}- or -SO₂NR^{8c}- where R⁸ is hydrogen atom or C₁₋₆ alkyl;

Y is a spacer having a main chain of 1 to 6 atoms;

provided that Ar is a ring of the formulae :



wherein symbols have the same meanings as defined above, and each ring may have substituents, when X is -SO₂NH-; and provided that Ar¹ is not biphenyl which may be substituted;

20 when X is -CONH- and Ar is any one of benzopyran, dihydrobenzopyran, dihydrobenzoxazine, dihydrobenzoxazole or tetrahydrobenzoxazepine;

WO 01/21577


PCT/JP00/06375

57

(excluding N-[2-(N,N-dimethylamino)methyl-6-tetralinyl]-4-biphenylcarboxamide);
namely compound of the formula (I') (excluding N-[2-(N,N-dimethylamino)methyl-6-tetralinyl]-4-biphenylcarboxamide) is a novel compound.

Preferred examples of compound of the formula (I') include compound of the formula (I'-1), (I'-2), (I'-3), (I'-4), (I'-5), (I'-6), (I'-7), (I'-8), (I'-9) or (I'-10).

In the above formulae (I'), (I'-1), (I'-2), (I'-3), (I'-4), (I'-5), (I'-6), (I'-7), (I'-8), (I'-9) and (I'-10), a ring of the formula :



wherein symbols have the same meanings as above, may have further 1 to 3 substituents at substitutable positions.

Examples of such substituents include "substituents" exemplified in the above Ar. Especially, preferred are formyl, optionally halogenated C₁₋₆ alkyl-carbonyl, optionally halogenated C₁₋₆ alkylsulfonyl, optionally halogenated C₁₋₆ alkyl (e.g., methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, hexyl, etc.), cyano, hydroxy, etc.

Examples of salts of compound (I) or (I') include salts with inorganic bases, ammonium salts, salts with organic bases, salts with inorganic acids, salts with organic acids, and salts with basic or acidic amino acids.

Preferred examples of salts with inorganic bases include alkali metal salts such as sodium salts and potassium salts; alkaline earth metal salts such as calcium salts, magnesium salts, barium salts; and aluminum salts.

Preferred examples of salts with organic bases include salts with trimethylamine, triethylamine, pyridine, picoline, ethanolamine, diethanolamine, triethanolamine,

WO 01/21577

PCT/JP00/06375

58

dicyclohexylamine, N,N-dibenzylethylenediamine.

Preferred examples of salts with inorganic acids include salts with hydrochloric acid, hydrobromic acid, nitric acid, sulfuric acid, phosphoric acid.

5 Preferred examples of salts with organic acids include salts with formic acid, acetic acid, trifluoroacetic acid, fumaric acid, oxalic acid, tartaric acid, maleic acid, citric acid, succinic acid, malic acid, methanesulfonic acid, benzenesulfonic acid, and p-toluenesulfonic acid,
10 3-chlorobenzoic acid.

Preferred examples of salts with basic amino acids include salts with arginine, lysine, ornithine. Preferred examples of salts with acidic amino acids include salts with aspartic acid, glutamic.

15 Among these salts, pharmaceutically acceptable salts are preferable. For instance, when compound (I) or (I') possesses an acidic functional group, it can form an inorganic salt such as an alkali metal salt (e.g., sodium salt, potassium salt, etc.), an alkaline earth metal salt
20 (e.g. calcium salt, magnesium salt, barium salt, etc.), and an ammonium salt. When compound (I) or (I') possesses a basic functional group, it can form an inorganic salt such as hydrochloride, sulfate, phosphate, hydrobromate, etc.; or an organic salt such as acetate, maleate, fumarate,
25 succinate, methanesulfonate, p-toluenesulfonate, citrate and tartrate.

Compounds (I) and (I') (hereinafter also abbreviated as a compound of the invention) can be either anhydrides or hydrates. A hydrate may have 0.5 to 3 water molecules.

30 In addition, a compound of the invention can be labeled using isotopes (e.g. ^3H , ^{14}C , and ^{35}S , etc.).

When a compound of the invention contains optical isomers, stereoisomers, regio isomers, rotational isomers,
35 these are included as a compound of the invention, and each of them can be obtained as a single substance by per se known

WO 01/21577

PCT/JP00/06375

59

synthesis methods and separation methods. For instance, when optical isomers exist in a compound of the invention, the optical isomers separated from the compound are included in a compound of the invention.

5 The optical isomers can be produced using per se known methods. Concretely, the optical isomer can be obtained by using an optically active synthetic intermediate, or subjecting the racemic mixture of the final product to optical resolution in accordance with common method.

10 Examples of optical resolution methods include per se known methods such as the fractional recrystallization method, chiral column method, diastereomer method, etc., which are described in detail below.

1) Fractional recrystallization method

15 The method which comprises allowing a racemate to form a salt with an optically active compound (e.g. (+)-mandelic acid, (-)-mandelic acid, (+)-tartaric acid, (-)-tartaric acid, (+)-1-phenethylamine, (-)-1-phenethylamine, cinchonine, (-)-cinchonidine, brucine, etc.), separating
20 the salt using a fractional recrystallization method, followed by, if desired, neutralizing process to obtain a free optical isomer.

2) Chiral column method

25 This method comprises subjecting a racemate or its salt to a column for separating an optical isomer (chiral column) for separation. For instance, in the case of liquid chromatography, an optical isomer mixture is added to the chiral column such as ENANTIO-OVM [produced by Tosoh] or CHIRAL series [produced by Daicel], which is developed
30 using water, various buffer solutions (e.g. phosphate buffer), organic solvents (e.g. ethanol, methanol, isopropanol, acetonitrile, trifluoroacetic acid, diethylamine, etc.) as single or mixed solutions, and the optical isomers are separated. Also, in the case of gas
35 chromatography, for instance, separation is conducted using a chiral column such as CP-Chirasil-DeX (produced by

WO 01/21577

PCT/JP00/06375

60

G.L.Science Co.).

3) Diastereomer method

In this method, a racemic mixture is subjected to a chemical reaction with an optically active reagent to give a diastereomer mixture, which is separated into a single substance by an ordinary separation means (e.g. fractional recrystallization, chromatography method, etc.). This single substance is subjected to removal of the optically active reagent part using chemical processing such as a hydrolysis reaction. For instance, when a compound of the invention possesses hydroxy or primary or secondary amino in its molecule, this compound is subjected to a condensation reaction with an optically active organic acid (e.g. MTPA [α -methoxy- α -(trifluoromethyl)phenylacetic acid], (-)-menthoxyacetic acid, etc.), to give the diastereomer in an ester form or an amide form, respectively. On the other hand, when a compound of the invention possesses carboxylic acid group, this compound is subjected to a condensation reaction with an optically active amine or alcohol reagent, to give the diastereomer in an amide form or an ester form, respectively. The separated diastereomer can be converted to an optical isomer of the original compound, by applying acidic hydrolysis or basic hydrolysis.

A prodrug of compound (I') is a compound which is converted to compound (I') by reactions involving enzymes and gastric acid, etc. under physiological conditions in the living body; in other words, a compound that is changed into compound (I') by enzymatically-caused oxidation, reduction and hydrolysis, and a compound that is changed into compound (I') by hydrolysis caused by gastric acid.

Examples of the prodrugs of compound (I') include compounds in which amino groups of compound (I') have been acylated, alkylated, or phosphorylated [e.g. compounds in which amino groups of compound (I') have been eicosanoylated, aranylated, pentylaminocarbonylated,

(5-methyl-2-oxo-1,3-dioxolen-4-yl)methoxycarbonylated, tetrahydrofuranylated, pyrrolidylmethylated, pivaloyloxymethylated, tert-butylated, etc.]; compounds in which hydroxyl groups of compound (I') have been
 5 acylated, alkylated, phosphorylated, borated (e.g. compounds in which hydroxyl groups of compound (I') have been acetylated, palmitoylated, propanoylated, pivaloylated, succinylated, fumarilated, alanilated, dimethylaminomethylcarbonylated, etc.); compounds in
 10 which carboxyl groups of compound (I') have been esterified or amidated [e.g. compounds in which carboxyl groups of compound (I') have been ethylesterified, phenylesterified, carboxylmethylesterified, dimethylaminomethylesterified,
 15 pivaloyloxymethylesterified, ethoxycarbonyloxyethylesterified, phthalidylesterified, (5-methyl-2-oxo-1,3-dioxolen-4-yl)methylesterified, cyclohexyloxycarbonylethylesterified, or methylamidated, etc.]. These compounds can be produced from compound (I')
 20 using per se known methods.

Also, a prodrug of compound (I') can be a compound which is changed to compound (I') by physiological conditions, as described in pages 163 to 198 of Molecular Design, Volume 7, "Development of Drugs," published in
 25 1990 by Hirokawa Shoten.

A compound of the invention can be produced in accordance with per se known methods such as methods described in WO9838156, WO9532967, and EP-A533266, etc.,
 30 or analogous methods thereto.

For instance, a compound of the invention can be produced in accordance with [Production method 1] to [Production method 6] which are described in detail below, or analogous methods thereto.

35 Compounds (II) to (XI) used as raw materials, can be used in the form of salts. As such salts, those exemplified

WO 01/21577

PCT/JP00/06375

62


as salts of the above compound (I) or (I') can be used.

In the following [Production method 1] to [Production method 6], when an alkylation reaction, a hydrolysis reaction, an amination reaction, an esterification reaction, an amidation reaction, an esterification reaction, an etherification reaction, an oxidation reaction, a reduction reaction, etc. are carried out, these reactions are carried out in accordance with per se known methods. Examples of such methods include the methods described in Organic Functional Group Preparations, Second Edition, Academic Press, Inc., published in 1989; Comprehensive Organic Transformations, VCH Publishers Inc., published in 1989, etc.

[Production method 1]

Compound (Ia) having $-(CH_2)_{w3}CONR^{8a}(CH_2)_{w4}-$ for X in formula (I), is produced, for instance, by the following amidation reaction.

(Amidation reaction)



wherein R^{8a} is hydrogen atom or an optionally halogenated C_{1-6} alkyl; other symbols have the same meanings as defined above.

As the "optionally halogenated C_{1-6} alkyl", those exemplified as "substituents" in the above "cyclic group which may have substituents" can be used.

The "amidation reaction" includes the following

WO 01/21577

PCT/JP00/06375

63

"method using a dehydration and condensation agent" and
"method using a reactive derivative of carboxylic acid".

i) Method using a dehydration and condensation agent

5 Compound (III), 1 to 5 equivalents of compound (II),
and 1 to 2 equivalents of a dehydration and condensation
agent are reacted in an inert solvent. If necessary, the
reaction can be carried out with the coexistence of 1 to
1.5 equivalents of 1-hydroxybenzotriazole (HOBT) and (or)
10 catalytic quantity to 5 equivalents of a base.

Examples of the "dehydrating and condensation agent"
include dicyclohexylcarbodiimide (DCC), 1-ethyl-3-(3-
dimethylaminopropyl)carbodiimide hydrochloride (WSC).
WSC is particularly preferable.

15 Examples of the "inert solvent" include nitrile
solvents (preferably acetonitrile), amide solvents
(preferably DMF), halogenated hydrocarbon solvents
(preferably dichloromethane), ether solvents (preferably
THF). Two or more kinds of these can be mixed in an
20 appropriate ratio for use.

Examples of the "base" include

1) for instance, strong bases such as hydrides of
alkali metals or alkaline earth metals (e.g. lithium
25 hydride, sodium hydride, potassium hydride, calcium
hydride, etc.), amides of alkali metals or alkaline earth
metals (e.g. lithium amide, sodium amide, lithium
diisopropylamide, lithium dicyclohexylamide, lithium
hexamethyldisilazide, sodium hexamethyldisilazide,
30 potassium hexamethyldisilazide, etc.), lower alkoxides of
alkali metals or alkaline earth metals (e.g. sodium
methoxide, sodium ethoxide, potassium tert-butoxide,
etc.);

2) for instance, inorganic bases such as hydroxides
35 of alkali metals or alkaline earth metals (e.g. sodium
hydroxide, potassium hydroxide, lithium hydroxide, barium

hydroxide, etc.), carbonates of alkali metals or alkaline earth metals (e.g. sodium carbonate, potassium carbonate, cesium carbonate, etc.) and hydrogencarbonates of alkali metals or alkaline earth metals (e.g. sodium hydrogencarbonate, potassium hydrogencarbonate, etc.); and

3) for instance, amines such as triethylamine, diisopropylethylamine, N-methylmorpholine, dimethylaminopyridine, DBU (1,8-diazabicyclo[5.4.0]undec-7-en), DBN (1,5-diazabicyclo[4.3.0]non-5-en); for instance, organic bases such as basic heterocyclic compounds of pyridine, imidazole, 2,6-lutidine, etc.

Among the above bases, triethylamine, 4-dimethylaminopyridine, etc., are preferable.

Reaction temperature is usually room temperature (0°C to 30°C, hereafter the same). Reaction time is, for instance, 10 to 24 hours.

ii) Method using a reactive derivative of carboxylic acid
A reactive derivative of compound (II) and 1 to 5 equivalents (preferably 1 to 3 equivalents) of compound (III) are reacted in an inert solvent. If necessary, the reaction can be carried out with the coexistence of 1 to 10 equivalents, preferably 1 to 3 equivalents of a base.

Examples of the "reactive derivative" of compound (II) include acid halides (e.g., acid chloride, acid bromide, etc.), mixed acid anhydrides (e.g. acid anhydrides with C₁₋₆ alkyl-carboxylic acid, C₆₋₁₀ aryl-carboxylic acid or C₁₋₆ alkylcarbonate), active esters (e.g. esters with phenol which may have substituents, 1-hydroxybenzotriazole or N-hydroxysuccinimide, etc.).

Examples of the "substituents" in the "phenol which may have substituents" include halogen atom (e.g. fluorine, chlorine, bromine, iodine, etc.), nitro, optionally halogenated C₁₋₆ alkyl, optionally halogenated C₁₋₆ alkoxy. The number of substituents is, for instance, 1 to 5.

As the "optionally halogenated C₁₋₆ alkyl" and "optionally halogenated C₁₋₆ alkoxy", those exemplified as "substituents" in the above "cyclic group which may have substituents" can be used.

5 Concrete examples of "phenol which may have substituents" include phenol, pentachlorophenol, pentafluorophenol, p-nitrophenol. The reactive derivative is, preferably, an acid halide.

10 Examples of the "inert solvent" include ether solvents, halogenated hydrocarbon solvents, aromatic solvents, nitrile solvents, amide solvents, ketone solvents, sulfoxide solvents, and water. Two or more kinds of these can be mixed in an appropriate ratio for use.

15 Especially, acetonitrile, THF, dichloromethane, chloroform, etc. are preferable.

As the "base", the same as above are used. The base is preferably sodium hydride, potassium carbonate, sodium carbonate, sodium hydroxide, potassium hydroxide, sodium hydrogencarbonate, potassium hydrogencarbonate, 20 triethylamine, pyridine, etc.

Reaction temperature is usually -20°C to 50°C, preferably room temperature. Reaction time is usually 5 minutes to 40 hours, preferably 1 to 18 hours.

Compound (III) can be produced by per se known methods. 25 For instance, 6-amino-2-(N,N-dimethylamino)methyltetraline or its salt can be produced in accordance with the methods described in WO9838156.

Also, 6-amino-2,3-dihydro-1-(2-dimethylaminoethyl)-1H-indole, 6-amino-3,4-dihydro-4-(2-dimethylaminoethyl)- 30 2H-1,4-benzoxazine, etc., can be produced in accordance with the methods described in WO9532967.

The above "method using a reactive derivative of carboxylic acid" can be also adopted when producing a corresponding sulfonamide derivative or sulfinamide 35 derivative, from the sulfonic acid of the formula :
Ar¹-(CH₂)_{w3}-SO₂OH (symbols have the same meanings as defined

WO 01/21577


PCT/JP00/06375

66

above), or the sulfinic acid of the formula : $\text{Ar}^1-(\text{CH}_2)_{w3}-\text{SOOH}$
(symbols have the same meanings as defined above).

[Production method 2]

- 5 Compound (Ib) having $-(\text{CH}_2)_{w3}-\text{COO}(\text{CH}_2)_{w4}-$ for X in the
formula (I), can be produced by the following
esterification reaction.
(Esterification reaction)



- 10 wherein symbols have the same meanings as defined above.

- A reactive derivative of compound (II) and 1 to 5
equivalents (preferably 1 to 3 equivalents) of compound
(IV) is reacted in an inert solvent. Usually, this reaction
is carried out with the coexistence of 1 to 10 equivalents,
15 preferably 1 to 3 equivalents of a base.

As the reactive derivative of compound (II), the same
as above is used. Especially, an acid halide is preferable.

- 20 Examples of the "inert solvent" include ether
solvents, halogenated hydrocarbon solvents, aromatic
solvents, nitrile solvents, amide solvents, ketone
solvents, sulfoxide solvents. Two or more kinds of these
can be mixed in an appropriate ratio for use. Especially,
acetonitrile, dichloromethane, chloroform, etc. are
preferable.

- 25 As the "base", the same one as above can be used. The
base is preferably sodium hydride, potassium carbonate,
sodium carbonate, sodium hydroxide, potassium hydroxide,
sodium hydrogencarbonate, potassium hydrogencarbonate,

WO 01/21577

PCT/JP00/06375

67

triethylamine, pyridine, etc.


Reaction temperature is usually -20°C to 50°C, preferably room temperature. Reaction time is usually 5 minutes to 40 hours, preferably 1 to 18 hours.

5

[Production method 3]

Compound (Ic) having $-(CH_2)_{w1}O(CH_2)_{w2}-$ for Y in the formula (I), can be produced by, for instance, the following etherification reaction.

10 (Etherification reaction)



wherein L is a leaving group, and other symbols have the same meanings as defined above.

15 Examples of the "leaving group" for L include halogen atom (e.g. chlorine, bromine, iodine, etc.), optionally halogenated C_{1-6} alkylsulfonyloxy (e.g. methanesulfonyloxy, ethanesulfonyloxy, trifluoromethanesulfonyloxy, etc.), C_{6-10} arylsulfonyloxy which may have substituents, hydroxy.

20 Examples of the "substituents" in the " C_{6-10} arylsulfonyloxy which may have substituents" include halogen atom (e.g. chlorine, bromine, iodine, etc.), optionally halogenated C_{1-6} alkyl, C_{1-6} alkoxy. The number of substituents is, for instance, 1 to 3. Concrete examples
25 of the C_{6-10} arylsulfonyloxy which may have substituents" include benzenesulfonyloxy, p-toluenesulfonyloxy, 1-naphthalenesulfonyloxy, 2-naphthalenesulfonyloxy.

The "leaving group" is preferably halogen atom (e.g.

WO 01/21577

PCT/JP00/06375

68

chlorine, bromine, iodine, etc.), methanesulfonyloxy, trifluoromethanesulfonyloxy, p-toluenesulfonyloxy.

Compound (IV') and about 1 to 5 equivalents
5 (preferably 1 to 2 equivalents) of compound (V) are reacted in inert solvent, with the coexistence of base.

As the "base", the same one as above can be used. The base is preferably potassium carbonate, sodium hydrogencarbonate, triethylamine, N-methylmorpholine,
10 pyridine, etc. The amount of the base used is usually about 1 to 5 equivalents relative to compound (V).

Examples of the "inert solvent" include alcohol solvents, ether solvents, halogenated hydrocarbon solvents, aromatic solvents, nitrile solvents, amide
15 solvents, ketone solvents, sulfoxide solvents, water. Two or more kinds of these can be mixed in an appropriate ratio for use. Especially, acetonitrile, N,N-dimethylformamide (DMF), acetone, ethanol, pyridine, etc., are preferable.

Reaction temperature is about -20°C to 100°C,
20 preferably room temperature to 80°C. Reaction time is, for instance, 5 hours to 1 day.

In the above production method, when the leaving group is hydroxy, Mitsunobu reaction can usually be used. In the Mitsunobu reaction, compound (V) and 0.5 to 5 equivalents
25 (preferably 1 to 1.5 equivalents) of compound (IV') are reacted in inert solvent with the coexistence of 0.5 to 5 equivalents (preferably 1 to 1.5 equivalents) of ethyl acetyldicarboxylate.

Examples of the inert solvent include ether solvents,
30 halogenated hydrocarbon solvents, aromatic solvents, nitrile solvents, amide solvents, ketone solvents, sulfoxide solvents. Two or more kinds of these can be mixed in an appropriate ratio for use. Especially, acetonitrile, dichloromethane, chloroform, etc. are
35 preferable.

Reaction temperature is usually -20°C to 50°C,

WO 01/21577

PCT/JP00/06375

69

preferably room temperature. Reaction time is usually 5 minutes to 40 hours, preferably 1 to 18 hours.


Compound (IV') can be produced by per se known methods.

For instance, 3-(N,N-dimethylamino)methyl-1,2,3,4-tetrahydro-7-quinolinol, 2-(N,N-dimethylamino)methyl-6-hydroxytetralin, 6-hydroxy-2-piperidinomethyltetralin, 2-[2-(N,N-dimethylamino)ethyl]-6-hydroxytetralin, 2-(N,N-dimethylamino)methyl-7-hydroxytetralin, 6-hydroxy-2-(N-methylamino)methyltetralin, etc., can be produced in accordance with the methods described in WO9838156.

[Production method 4]

Compound (Id) having $-(CH_2)_{w3}NR^{8a}CO(CN_2)_{w4}-$ for X in the formula (I), can be produced, for instance, by the following amidation reaction.

(Amidation reaction)



wherein symbols have the same meanings as defined above.

This Production method is carried out in accordance with the above Production method 1.

[Production method 5]


Compound (Ie) having $-(CH_2)_{w5}NHCONR^{8a}(CN_2)_{w6}-$ for X in the formula (I), can be produced, for instance, by the following urea reaction.

(Urea reaction)

WO 01/21577

PCT/JP00/06375

70



wherein symbols have the same meanings as defined above.

Compound (IX) and 1 to 5 equivalents (preferably 1 to 5 or 1.5 equivalents) of compound (VIII) is reacted in an inert solvent with the coexistence of a base.

As the "base", the same one as above can be used. The base is preferably potassium carbonate, sodium carbonate, sodium hydroxide, potassium hydroxide, sodium hydrogencarbonate, potassium hydrogencarbonate, triethylamine, pyridine, etc.

Examples of the "inert solvent" include alcohol solvents, ether solvents, halogenated hydrocarbon solvents, aromatic solvents, nitrile solvents, amide solvents, ketone solvents, sulfoxide solvents, water. Two or more kinds of these can be mixed in an appropriate ratio for use. Especially, acetonitrile, DMF, acetone, ethanol, pyridine, etc. are preferable.

Reaction temperature is usually -20°C to 100°C, preferably room temperature to 80°C. Reaction time is, for instance, 0.5 hour to 1 day.

[Production method 6]

Compound (If) having, for Ar¹, a ring assembly aromatic


WO 01/21577

PCT/JP00/06375

71

group ($\text{Ar}^2\text{-Ar}^3$) which may have substituents in the formula (I), can be produced by, for instance, the following aryl-coupling reaction.

(Aryl-coupling reaction)



5

wherein Ar^2 and Ar^3 are monocyclic aromatic groups or condensed aromatic groups, each of which may have substituents; L^1 is hydroxy or C_{1-6} alkyl; L^2 is halogen (preferably chlorine, bromine) or

10 trifluoromethanesulfonyloxy; other symbols have the same meanings as defined above.

As "substituents", "monocyclic aromatic groups" and "condensed aromatic groups" in the "monocyclic aromatic groups or condensed aromatic groups, each of which may have substituents" for Ar^2 and Ar^3 , those exemplified as the above Ar^1 can be used. Especially, it is preferable that both of Ar^2 and Ar^3 are phenyl groups which may have substituents, and $\text{Ar}^2\text{-Ar}^3$ is biphenyl which may have substituents.

20 The aryl-coupling reaction can be carried out in accordance with per se known methods such as the method described in Acta. Chemica Scandinavia, pp. 221-230, 1993, or methods analogous thereto.

Compound (X) and 1 to 3 equivalents (preferably 1 to 25 1.5 equivalents) of compound (XI) are reacted in an inert solvent in the presence of a base and a transition metal catalyst.

As the base, the same one as above can be used. The

WO 01/21577

PCT/JP00/06375

72

base is preferably sodium carbonate, sodium hydrogencarbonate, etc.

The amount of the "base" used is, for instance, about 1 to 10 equivalents relative to compound (XI).

5 Examples of the "transition metal catalyst" include palladium catalyst, nickel catalyst. Examples of the "palladium catalyst" include tetrakis(triphenylphosphine)palladium (O), palladium acetate, bis (triphenylphosphine) palladium (II) chloride,
10 palladium-carbon. Examples of the "nickel catalyst" include tetrakis(triphenylphosphine) nickel (O).

The amount of the "transition metal catalyst" used is about 0.01 to 1 equivalent, preferably about 0.01 to 0.5 equivalent, relative to compound (XI).

15 Reaction temperature is room temperature to 150°C, preferably about 80°C to 150°C. Reaction time is, for instance, about 1 to 48 hours.

Examples of the "inert solvent" include water, alcohol solvents, aromatic solvents. Two or more kinds of these
20 can be mixed in an appropriate ratio for use. Especially, a single solvent such as water, ethanol and toluene; or a mixed solvent of two or more kinds of these is preferable.

Examples of the above "alcohol solvents" include methanol, ethanol, isopropanol, tert-butanol.

25 Examples of the above "ether solvents" include diethylether, tetrahydrofuran (THF), 1,4-dioxane, 1,2-dimethoxyethane.

Examples of the above "halogenated hydrocarbon solvents" include dichloromethane, chloroform, 1,2-
30 dichloroethane, carbon tetrachloride.

Examples of the above "aromatic solvents" include benzene, toluene, xylene, pyridine.

Examples of the above "hydrocarbon solvents" include hexane, pentane, cyclohexane.

35 Examples of the above "amide solvents" include N,N-dimethylformamide (DMF), N,N-dimethylacetamide, N-

WO 01/21577

PCT/JP00/06375

73

methylpyrrolidone.

Examples of the above "ketone solventd" include acetone, methylethylketone.

5 Examples of the above "sulfoxide solvents" include dimethylsulfoxide (DMSO).

Examples of the above "nitrile solvents" include acetonitrile, propionitrile.

10 In a compound of the invention thus obtained, the intramolecular functional group can be converted to a desired functional group by combining per se known chemical reactions. Examples of the chemical reactions include oxidation reaction, reduction reaction, alkylation reaction, hydrolysis reaction, amination reaction, esterification reaction, aryl-coupling reaction,
15 deprotection reaction.

In each of the above reactions, when the raw material compounds possess amino, carboxy, hydroxy, and/or carbonyl as substituents, protecting groups which are generally used in peptide chemicals, etc., can be introduced into these
20 groups, and the desired compound can be obtained by removing the protecting groups after the reaction if necessary.

Examples of the protecting group for amino include formyl, C₁₋₆ alkyl-carbonyl (e.g. acetyl, propionyl, etc.), C₁₋₆ alkoxy-carbonyl (e.g. methoxycarbonyl,
25 ethoxycarbonyl, tert-butoxycarbonyl, etc.), benzoyl, C₇₋₁₀ aralkyl-carbonyl (e.g. benzylcarbonyl, etc.), C₇₋₁₄ aralkyloxy-carbonyl (e.g. benzyloxycarbonyl, 9-fluorenylmethoxycarbonyl, etc.), trityl, phthaloyl, N,N-dimethylaminomethylene, silyl (e.g. trimethylsilyl, triethylsilyl, dimethylphenylsilyl, tert-
30 butyldimethylsilyl, tert-butyldiethylsilyl, etc.), C₂₋₆ alkenyl (e.g. 1-allyl, etc.) . These groups may be substituted by 1 to 3 of halogen atom (e.g. fluorine, chlorine, bromine, iodine, etc.), C₁₋₆ alkoxy (e.g. methoxy, ethoxy, propoxy, etc.) or nitro, etc.
35

Examples of the protecting group for carboxy include

C₁₋₆ alkyl (e.g. methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, etc.), C₇₋₁₁ aralkyl (e.g. benzyl, etc.), phenyl, trityl, silyl (e.g. trimethylsilyl, triethylsilyl, dimethylphenylsilyl, tert-butyldimethylsilyl, tert-
5 butyldiethylsilyl, etc.), C₂₋₆ alkenyl (e.g. 1-allyl, etc.). These groups may be substituted by 1 to 3 of halogen atom (e.g. fluorine, chlorine, bromine, iodine, etc.), C₁₋₆ alkoxy (e.g. methoxy, ethoxy, propoxy, etc.) or nitro.

Examples of the protective group for hydroxy include
10 C₁₋₆ alkyl (e.g. methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, etc.), phenyl, trityl, C₇₋₁₀ aralkyl (e.g. benzyl, etc.), formyl, C₁₋₆ alkyl-carbonyl (e.g. acetyl, propionyl, etc.), benzoyl, C₇₋₁₀ aralkyl-carbonyl (e.g. benzylcarbonyl, etc.), 2-tetrahydropyranyl, 2-
15 tetrahydrofuranyl, silyl (e.g. trimethylsilyl, triethylsilyl, dimethylphenylsilyl, tert-butyldimethylsilyl, tert-butyldiethylsilyl, etc.), C₂₋₆ alkenyl (e.g. 1-allyl, etc.). These groups may be substituted by 1 to 3 of halogen atom (e.g. fluorine,
20 chlorine, bromine, iodine, etc.), C₁₋₆ alkyl (e.g. methyl, ethyl, n-propyl, etc.), C₁₋₆ alkoxy (e.g. methoxy, ethoxy, propoxy, etc.) or nitro, etc. can be substituted for these groups.

Examples of the protecting group for carbonyl include
25 cyclic acetal (e.g. 1,3-dioxane, etc.), and non-cyclic acetal (e.g. di-C₁₋₆ alkylacetal, etc.).

Removal of the above protecting groups can be carried out in accordance with per se known methods such as those described in Protective Groups in Organic Synthesis,
30 published by John Wiley and Sons (1980). For instance, the methods using acid, base, ultraviolet light, hydrazine, phenylhydrazine, sodium N-methyldithiocarbamate, tetrabutylammonium fluoride, palladium acetate, trialkylsilyl halide (e.g. trimethylsilyl iodide,
35 trimethylsilyl bromide, etc.), and a reduction method, etc. can be used.

WO 01/21577

PCT/JP00/06375

75

A compound of the invention can be isolated and purified by per se known methods such as solvent extraction, changing of liquid properties, transdissolution, crystallization, recrystallization, chromatography, etc.

5 It is also possible to isolate and purify the raw material compounds of a compound of the invention, or their salts using the same known methods as above, but they can also be used as raw materials in the next process as a reaction mixture without being isolated.

10

A compound of the invention possesses an excellent MCH receptor antagonistic action, therefore, it is useful as an agent for preventing or treating diseases caused by MCH.

15 Also, a compound of the invention is low in toxicity, and is excellent in oral absorbency and intracerebral transitivity.

Therefore, a melanin-concentrating hormone antagonist (hereafter, also abbreviated as "MCH antagonist") comprising a compound of the invention can be
20 safely administered to mammals (e.g. rats, mice, guinea pigs, rabbits, sheep, horses, swine, cattle, monkeys, humans, etc.) as an agent for preventing or treating diseases caused by MCH.

Here, examples of the diseases caused by MCH include
25 obesity (e.g. malignant mastocytosis, exogenous obesity, hyperinsulinar obesity, hyperplasmic obesity, hypophyseal adiposity, hypoplasmic obesity, hypothyroid obesity, hypothalamic obesity, symptomatic obesity, infantile obesity, upper body obesity, alimentary obesity,
30 hypogonadal obesity, systemic mastocytosis, simple obesity, central obesity, etc.), hyperphagia, emotional disorders, reproductive function disorders, memory disorders, dementia, hormonal disorders.

A compound of the invention is also useful as an agent
35 for preventing or treating lifestyle diseases such as diabetes, diabetic complications (e.g. diabetic

WO 01/21577

PCT/JP00/06375

76

retinopathy, diabetic neuropathy, diabetic nephropathy, etc.), arteriosclerosis, and gonitis.

Further, a compound of the invention is useful as an anorectic agent.

5 A MCH antagonist and a pharmaceutical composition of the invention can be used in combination with an alimentary therapy (e.g., alimentary therapy for diabetes) and exercise.

10 A MCH antagonist and a pharmaceutical composition of the invention can be produced by subjecting compound (I) or compound (I') respectively, as it is, or together with a pharmacologically acceptable carrier, to pharmaceutical manufacturing process in accordance with a per se known means.

15 Here, examples of the pharmacologically acceptable carriers include various organic or inorganic carrier substances which are commonly used as materials for pharmaceutical preparations, such as excipients, lubricants, binders, and disintegrators in solid
20 preparations; solvents, solubilizing agents, suspending agents, isotonizing agents, buffering agents, soothing agents, in liquid preparations. Also, in the pharmaceutical manufacturing process, additives such as antiseptics, antioxidants, coloring agents, sweeteners,
25 absorbents, moistening agents, can be used, if necessary.

Examples of the excipients include lactose, sucrose, D-mannitol, starch, cornstarch, crystalline cellulose, light anhydrous silicic acid.

30 Examples of the lubricants include magnesium stearate, calcium stearate, talc, colloidal silica.

Examples of the binders include crystalline cellulose, sucrose, D-mannitol, dextrin, hydroxypropylcellulose, hydroxypropylmethylcellulose, polyvinylpyrrolidone, starch, saccharose, gelatin,
35 methylcellulose, carboxymethylcellulose sodium.

Examples of the disintegrators include starch,

WO 01/21577

PCT/JP00/06375

77

carboxymethylcellulose, carboxymethylcellulose calcium, crosscarmellose sodium, carboxymethylstarch sodium, low-substituted hydroxypropylcellulose (L-HPC).

5 Examples of the solvents include distilled water for injection, alcohol, propylene glycol, macrogol, sesame oil, corn oil.

10 Examples of the solubilizing agents include polyethylene glycol, propylene glycol, D-mannitol, benzyl benzoate, ethanol, trisaminomethane, cholesterol, triethanolamine, sodium carbonate, sodium citrate.

15 Examples of the suspending agents include surfactants such as stearyltriethanolamine, sodium lauryl sulfate, lauryl amino propionic acid, lecithin, benzalkonium chloride, benzethonium chloride, glyceryl monostearate; or hydrophilic polymers such as polyvinyl alcohol, polyvinylpyrrolidone, carboxymethylcellulose sodium, methylcellulose, hydroxymethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose.

20 Examples of the isotonizing agents include glucose, D-sorbitol, sodium chloride, glycerin, D-mannitol.

Examples of the buffering agents include buffer solutions of phosphate, acetate, carbonate and citrate.

Examples of the soothing agents include benzyl alcohol.

25 Examples of the antiseptics include paraoxybenzoates, chlorobutanol, benzyl alcohol, phenethylalcohol, dehydroacetic acid, and sorbic acid.

Examples of the antioxidants include sulfite, ascorbic acid.

30 A MCH antagonist and a pharmaceutical composition of the invention can be safely administered orally or parenterally (e.g. by local, rectal and intravenous administration) in various dosage forms, for instance, as oral drugs such as tablets (including sugar-coated tablets and film-coated tablets), powders, granules, capsules
35 (including soft capsules), solutions; and parenteral

preparations such as injections (e.g. subcutaneous injections, intravenous injections, intramuscular injections, intraperitoneal injections, etc.), external preparations (e.g. nasal preparations, percutaneous preparations, ointments, etc.), suppositories (e.g. rectal suppositories, vaginal suppositories, etc.), sustained-release preparations (e.g. sustained-release microcapsules, etc.), pellets, drip infusions, etc.

The content of compound (I) in a MCH antagonist of the invention and the content of compound (I') in a pharmaceutical composition of the invention are, for instance, about 0.1 to 100 weight percent of the MCH antagonist or whole pharmaceutical composition, respectively.

The dose of a MCH antagonist and a pharmaceutical composition of the invention can be appropriately selected depending on the subject of administration, route of administration, disease, etc.

For instance, the dose per day when a MCH antagonist or a pharmaceutical composition of the invention is orally administered to an adult obesity patient (body weight: about 60 kg), is about 0.1 to about 500 mg, preferably about 1 to about 100 mg, more preferably about 5 to about 100 mg, in terms of compound (I) or compound (I'), each of which is an active ingredient. These amounts can be divided into one to several doses per day for administration.

The MCH antagonist and pharmaceutical composition of the invention can be used in combination with other concomitant drugs which do not interfere with the MCH antagonist and pharmaceutical composition of the invention, for the purpose of "strengthening of therapeutic effect against obesity", "reduction of dose of MCH antagonist", etc. Examples of the concomitant drugs include a "agents for treating diabetes", "agents for treating diabetic complications", "agents for treating obesity other than MCH antagonists", "agents for treating

WO 01/21577

PCT/JP00/06375

79

hypertension", "agents for treating hyperlipidemia (agents for treating arteriosclerosis)", "agents for treating arthritis", "antianxiety agents", "antidepressant". Two or more kinds of these concomitant drugs can be combined
5 in an appropriate ratio for use.

Examples of the above "agents for treating diabetes" include insulin sensitizers, insulin secretion enhancers, biguanides, insulins, α -glucosidase inhibitors, β 3 adrenaline receptor agonists.

10 Examples of the insulin sensitizers include pioglitazone or its salt (preferably hydrochloride), troglitazone, rosiglitazone or its salt (preferably maleate), JTT-501, GI-262570, MCC-555, YM-440, DRF-2593, BM-13-1258, KRP-297, R-119702.

15 Examples of the insulin secretion enhancers include sulfonylureas. Concrete examples of the sulfonylureas include tolbutamide, chlorpropamide, trazamide, acetohexamide, glycopyramide and its ammonium salt, glibenclamide, gliclazide, glimepiride.

20 Other than the above, examples of insulin secretion enhancers include repaglinide, nateglinide, mitiglinide (KAD-1229), JTT-608.

Examples of biguanides include metformin, buformin, phenformin.

25 Examples of insulins include animal insulins extracted from bovine or porcine pancreas; semi-synthetic human insulin which is enzymatically synthesized from insulin extracted from porcine pancreas; human insulin synthesized by genetic engineering, using Escherichi Coli and yeast. As insulin, also employed are insulin-zinc
30 containing 0.45 to 0.9 (w/w)% of zinc; protamine-insulin-zinc produced from zinc chloride, protamine sulfate and insulin. In addition, insulin can be an insulin fragment or derivative (e.g. INS-1, etc.).

35 Insulin can also include various types such as ultra immediate action type, immediate action type, two-phase

WO 01/21577

PCT/JP00/06375

80

type, intermediate type, prolonged action type, etc., and these can be selected depending on the pathological conditions of patients.

5 Examples of α -glucosidase inhibitors include acarbose, voglibose, miglitol, emiglitate.

Examples of β 3 adrenaline receptor agonists include AJ-9677, BMS-196085, SB-226552, AZ40140.

10 Other than the above, examples of the "agents for treating diabetes" include ergoset, pramlintide, leptin, BAY-27-9955.

Examples of the above "agents for treating diabetic complications" include aldose reductase inhibitors, glycation inhibitors, protein kinase C inhibitors.

15 Examples of aldose reductase inhibitors include torulestat; eparlestat; imirestat; zenarestat; SNK-860; zopolrestat; ARI-509; AS-3201.

Examples of glycation inhibitors include pimagedine.

Examples of protein kinase C inhibitors include NGF, LY-333531.

20 Other than the above, examples of "agents for treating diabetic complications" include alprostadil, thiapride hydrochloride, cilostazol, mexiletine hydrochloride, ethyl eicosapentate, memantine, pimagedline (ALT-711).

25 Examples of the above "agents for treating obesity other than MCH antagonists" include lipase inhibitors and anorectics.

Examples of lipase inhibitors include orlistat.

30 Examples of anorectics include mazindol, dexfenfluramine, fluoxetine, sibutramine, baiamine, (S)-sibutramine, SR-141716, NGD-95-1.

Other than the above, examples of "agents for treating obesity other than MCH antagonists" include lipstatin.

35 Examples of the above "agents for treating hypertension" include angiotensin converting enzyme inhibitors, calcium antagonists, potassium channel openers, angiotensin II antagonists,

WO 01/21577

PCT/JP00/06375

81

Examples of angiotensin converting enzyme inhibitors include captopril, enalapril, alacepril, delapril (hydrochloride), lisinopril, imidapril, benazepril, cilazapril, temocapril, trandolapril, manidipine
5 (hydrochloride).

Examples of calcium antagonists include nifedipine, amlodipine, efonidipine, nicardipine.

Examples of potassium channel openers include levcromakalim, L-27152, AL0671, NIP-121.

10 Examples of angiotensin II antagonists include losartan, candesartan cilexetil, valsartan, irbesartan, CS-866, E4177.

Examples of the above "agents for treating hyperlipidemia (agents for treating arteriosclerosis)"
15 include HMG-CoA reductase inhibitors, fibrate compounds.

Examples of HMG-CoA reductase inhibitors include pravastatin, simvastatin, lovastatin, atorvastatin, fluvastatin, lipantil, cerivastatin, itavastatin, ZD-4522, or their salts (e.g. sodium salts, etc.).

20 Examples of fibrate compounds include bezafibrate, clinofibrate, clofibrate, simfibrate.

Examples of the above "agents for treating arthritis" include ibuprofen.

25 Examples of the above "antianxiety agents" include chlordiazepoxide, diazepam, oxazolam, medazepam, cloxazolam, bromazepam, lorazepam, alprazolam, fludiazepam.

30 Examples of the above "antidepressants" include fluoxetine, fluvoxamine, imipramine, paroxetine, sertraline.

The timing of administration of the above concomitant drugs is not limited. The MCH antagonist or pharmaceutical composition and the concomitant drugs can be administered to the subject simultaneously or at staggered times.

35 The dosages of the concomitant drugs can be determined in accordance with clinically used dosages, and can be

appropriately selected according to the subject of administration, route of administration, diseases and combinations of drugs, etc.

The administration forms for the concomitant drugs are not particularly limited as long as a MCH antagonist or a pharmaceutical composition are used in combination with a concomitant drugs at the time of administration. Examples of such administration forms includes 1) administration of a single preparation obtained by simultaneous preparation of MCH antagonist or pharmaceutical composition together with concomitant drugs, 2) simultaneous administration of two kinds of preparations obtained by separate preparation of MCH antagonist or pharmaceutical composition, and concomitant drugs, through the same route of administration, 3) staggered administration of two kinds of preparations obtained by separate preparation of MCH antagonist or pharmaceutical composition, and concomitant drugs, through the same route of administration, 4) simultaneous administration of two kinds of preparations obtained by separate preparation of MCH antagonist or pharmaceutical composition, and concomitant drugs, through different routes of administration, 5) staggered administration of two kinds of preparations obtained by separate preparation of MCH antagonist or pharmaceutical composition, and concomitant drugs, through different routes of administration (for instance, administration of MCH antagonist or pharmaceutical composition; and concomitant drugs in this order; or administration in reverse order).

The ratio of combination of MCH antagonist or pharmaceutical composition with concomitant drugs can be appropriately selected in accordance with the subject of administration, route of administration and diseases, etc.

This invention further relates to "a pharmaceutical comprising a melanin-concentrating hormone antagonist in

combination with at least one species selected from the group consisting of an agent for treating diabetes, an agent for treating hypertension and an agent for treating arteriosclerosis".

5 Here, the "melanin-concentrating hormone antagonist" is not especially limited as long as it is a compound having a melanin-concentrating hormone antagonistic action, and may be either of a peptide compound or a non-peptide compound.

10 As "an agent for treating diabetes", "an agent for treating hypertension" and "an agent for treating arteriosclerosis", those exemplified as the above concomitant drugs can be mentioned.

15 These drugs can be used in the same manner as in the above "combination of MCH antagonist of the invention with concomitant drugs".

20 The pharmaceutical provides excellent effects such as "strengthening of therapeutic effect against obesity", "reduction of dose of MCH antagonist", etc. as compared to single use of each drug.

BEST MODE FOR CARRYING OUT THE INVENTION

25 This invention will be explained further in detail by the following Reference Examples, Examples, Preparation Examples, and Experimental Examples. However, these do not limit this invention, and they can be changed within the scope that does not deviate from the scope of this invention.

30 In the following Reference Examples and Examples, "room temperature" means 0 to 30°C. Anhydrous magnesium sulfate or anhydrous sodium sulfate was used to dry the organic layer. "%" means percent by weight, unless otherwise specified.

35 Infrared absorption spectra were determined by the diffuse reflectance method, using fourier transform type infrared spectrophotometer.

WO 01/21577

PCT/JP00/06375

84

FABMS (pos) is mass spectrum determined by the (+) method, in Fast Atom Bombardment Mass Spectrometry.

Other symbols used in the description have the
5 following meanings.

	s	: singlet
	d	: doublet
	t	: triplet
	q	: quartet
10	m	: multiplet
	br	: broad
	J	: coupling constant
	Hz	: Hertz
	CDCl ₃	: heavy chloroform
15	DMSO-d ₆	: heavy dimethylsulfoxide
	THF	: tetrahydrofuran
	DMF	: N,N-dimethylformamide
	DMSO	: dimethylsulfoxide
	WSCD	: 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide
20	WSC	: 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide hydrochloride
	¹ H-NMR	: proton nuclear resonance (Free substances were usually measured in 25 CDCl ₃ .)
	IR	: infrared absorption spectrum
	Me	: methyl
	Et	: ethyl
	HOBt	: 1-hydroxy-1H-benzotriazole
30	IPE	: diisopropyl ether
	DMAP	: 4-dimethylaminopyridine

In this specification and drawings, when bases and
35 amino acids are shown by codes, these codes are based on those by the IUPAC-IUB Commission on Biochemical

WO 01/21577

PCT/JP00/06375

85

Nomenclature or common codes in the concerned fields.
Examples of these codes are shown below. Also, where some
optical isomers of amino acids can exist, the L form is shown
unless otherwise specified.

5	DNA	: deoxyribonucleic acid
	cDNA	: complementary deoxyribonucleic acid
	A	: adenine
	T	: thymine
	G	: guanine
10	C	: cytosine
	RNA	: ribonucleic acid
	mRNA	: messenger ribonucleic acid
	dATP	: deoxyadenosine triphosphate
	dTTP	: deoxythymidine triphosphate
15	dGTP	: deoxyguanosine triphosphate
	dCTP	: deoxycytidine triphosphate
	ATP	: adenosine triphosphate
	EDTA	: ethylenediamine tetraacetic acid
	SDS	: sodium dodecyl sulfate
20	EIA	: enzyme immunoassay
	Gly	: glycine
	Ala	: alanine
	Val	: valine
	Leu	: leucine
25	Ile	: isoleucine
	Ser	: serine
	Thr	: threonine
	Cys	: cysteine
	Met	: methionine
30	Glu	: glutamic acid
	Asp	: aspartic acid
	Lys	: lysine
	Arg	: arginine
	His	: histidine
35	Phe	: phenylalanine
	Tyr	: tyrosine

WO 01/21577

PCT/JP00/06375

86

	Tro	: tryptophan
	Pro	: proline
	Asn	: asparagine
	Gln	: glutamine
5	pGl	: pyroglutamine
	Me	: methyl group
	Et	: ethyl group
	Bu	: butyl group
	Ph	: phenyl group
10	TC	: thiazolidine-4(R)-carboxamide group

Substituents, protecting groups and reagents frequently used in this specification, are shown by the following symbols.

15	Tos	: p-toluenesulfonyl
	CHO	: formyl
	Bzl	: benzyl
	Cl ₂ Bzl	: 2,6-dichlorobenzyl
	Bom	: benzyloxymethyl
20	Z	: benzyloxycarbonyl
	Cl-Z	: 2-chlorobenzyloxycarbonyl
	Br-Z	: 2-bromobenzyloxycarbonyl
	Boc	: t-butoxycarbonyl
	DNP	: dinitrophenol
25	Trt	: trityl
	Bom	: t-butoxymethyl
	Fmoc	: N-9-fluorenylmethoxycarbonyl
	HOBt	: 1-hydroxybenzotriazole
	HOObt	: 3,4-dihydro-3-hydroxy-4-oxo-1,2,3-
30		benzotriazine
	HONB	: 1-hydroxy-5-norbornene-2,3-
		dicarbodiimide
	DCC	: N,N'-dicyclohexylcarbodiimide

35 SEQ ID NO in the SEQUENCE LISTING in the specification of the present application shows the following sequences.

- [SEQ ID NO : 1] shows a synthetic DNA used for screening of cDNA coding rat SLC-1.
- [SEQ ID NO : 2] shows a synthetic DNA used for screening
5 of cDNA coding rat SLC-1.
- [SEQ ID NO : 3] shows an entire amino acid sequence of rat SLC-1.
- [SEQ ID NO : 4] shows an entire base sequence of rat SLC-1cDNA wherein Sal I recognition sequence was added to the 5' side,
10 and Spe I recognition sequence was added to the 3' side.
- [SEQ ID NO : 5] shows riboprobe used to determine the quantity of SLC-1mRNA expressed in each clone of rat SLC-1 expression CHO cells.
- [SEQ ID NO : 6] shows a synthetic DNA used to obtain cDNA
15 for coding of human SLC-1.
- [SEQ ID NO : 7] shows a primer used to make double-strand cDNA for coding human SLC-1.
- [SEQ ID NO : 8] shows an entire base sequence of cDNA for coding human SLC-1.
- 20 [SEQ ID NO : 9] shows an entire amino acid sequence of human SLC-1.
- [SEQ ID NO : 10] shows a synthetic DNA used for screening of cDNA for coding human SLC-1(S).
- [SEQ ID NO : 11] shows a synthetic DNA used for screening
25 of cDNA for coding human SLC-1(S).
- [SEQ ID NO : 12] shows a synthetic DNA used for screening of cDNA for coding human SLC-1(L).
- [SEQ ID NO : 13] shows a synthetic DNA used for screening
30 of cDNA for coding human SLC-1(L).
- [SEQ ID NO : 14] shows an entire base sequence of human SLC-1(S) cDNA wherein Sal I recognition sequence was added to the 5' side, and Spe I recognition sequence was added to the 3' side.
- 35 [SEQ ID NO : 15] shows an entire base sequence of human SLC-1(L) cDNA wherein Sal I recognition sequence was added

WO 01/21577

PCT/JP00/06375

88


to the 5' side, and Spe I recognition sequence was added to the 3' side.

[SEQ ID NO : 16] shows riboprobe used to determine the quantity of SLC-1mRNA expressed in each clone of human SLC-1(S) expression CHO cells and SLC-1(L) expression CHO cells.

Transformant Escherichia coli DH10B/phSLC1L8 transformed by plasmid containing DNA which codes the base sequence shown by SEQ ID NO : 9, obtained in Reference Example 1 - 6, is on deposit with National Institute of Bioscience and Human-Technology (NIBH), Agency of Industrial Science and Technology, Ministry of International Trade and Industry, as deposit number FERM BP-6632 from February 1, 1999; and with the Institute for Fermentation, Osaka, Japan (IFO), as deposit number IFO 16254 from January 21, 1999.

Reference Example 1

2-(R)-[2-(N,N-Dimethylamino)ethyl]-6-(4-[(4-methoxyphenyl)carbonyloxy]benzyloxy)tetralin



Diethyl azodicarboxylate (40% toluene solution, 0.95 g) was added dropwise to THF solution (6 ml) of 2-(R)-[2-(N,N-dimethylamino)ethyl]-6-hydroxytetralin (300 mg), 4-(hydroxymethyl)phenyl 4-methoxybenzoate (530 mg), and triphenylphosphine (430 mg) under ice-cooling. After stirring for 2 hours at room temperature, the reaction mixture was concentrated. The residue was purified using alumina column chromatography (development solvent; hexane ~ hexane : ethyl acetate = 10:1), and the titled compound

WO 01/21577

PCT/JP00/06375

89

(320 mg) was obtained after recrystallization (ethyl acetate-hexane).


Melting point: 111 - 114°C

$[\alpha]_D^{20} = +44.4^\circ$ (c = 0.502, methanol)

5

Reference Example 2

N-Phenyl-4-[[2-(2-piperidinoethyl)-6-tetralinyl]oxymethyl]benzamide



10 Triethylamine (0.11 ml) was added to THF suspension
(3 ml) of 4-[[2-(2-piperidinoethyl)-6-tetralinyl]oxymethyl]benzoate (300 mg). Further, THF
solution (0.5 ml) of trimethylacetyl chloride (92 mg) was
added dropwise under ice-cooling, which was stirred for 30
15 minutes. The temperature of the reaction mixture was
raised to room temperature, which was stirred for 1 hour.
THF solution (0.5 ml) of aniline (85 mg) was added dropwise
to the reaction mixture under ice-cooling, which was
stirred for 1 hour. After the reaction mixture was stirred
20 for 24 hours at room temperature, saturated sodium
bicarbonate solution was added, and extraction was
conducted using a mixed solution of ethyl acetate and THF.
The organic layer was washed with water and saturated
aqueous sodium chloride solution, dried, and then
25 concentrated. The residue was recrystallized from THF-
methanol-IPE to give the titled compound (150 mg).
Melting point: 183 - 185°C


Reference Example 3

30 4-[[2-(2-Piperidinoethyl)-6-tetralinyl]oxymethyl]-N-(2-
pyridinyl)benzamide

WO 01/21577

PCT/JP00/06375

90




Triethylamine (0.11 ml) was added to THF suspension (6 ml) of 4-[[2-(2-piperidinoethyl)-6-tetralinyl]oxymethyl]benzoate (300 mg). Trimethylacetyl chloride (0.095 ml) was added dropwise to the obtained suspension under ice-cooling, which was stirred for 30 minutes. The temperature of the reaction mixture was raised to room temperature, which was stirred for 1 hour.

THF solution (1.0 ml) of 2-aminopyridine (110 mg) was added dropwise to the reaction mixture under ice-cooling, which was stirred for 1 hour. Then the reaction mixture was stirred at room temperature for 6 hours, and at 60°C for 12 hours, which was refluxed with heating for 6 hours.

Saturated sodium bicarbonate solution was added to the reaction mixture, and extraction was conducted using ethyl acetate. The organic layer was washed with water and saturated aqueous sodium chloride solution, dried, and then concentrated. The residue was purified using alumina column chromatography (development solvent: THF), and recrystallized (ethyl acetate-IPE) to give the titled compound (30 mg).

Reference Example 4

4-[[2-(2-Piperidinoethyl)-6-tetralinyl]oxymethyl]-N-(2-quinolinyl)benzamide



WO 01/21577

PCT/JP00/06375


91

Triethylamine (0.22 ml) was added to THF suspension (6 ml) of 4-[[2-(2-piperidinoethyl)-6-tetralinyl]oxymethyl]benzoate (300 mg). Further, trimethylacetyl chloride (0.095 ml) was added dropwise to under ice-cooling, which was stirred for 30 minutes. The temperature of the reaction mixture was raised to room temperature, which was stirred for 1 hour. THF solution (1.0 ml) of 2-aminoquinoline (170 mg) was added dropwise to the reaction mixture under ice-cooling, which was stirred at room temperature for 12 hours. Saturated sodium bicarbonate solution was added to the reaction mixture, and extraction was conducted using ethyl acetate. The organic layer was washed with water and saturated aqueous sodium chloride solution, dried, and then concentrated. The residue was purified using alumina column chromatography (development solvent: THF), and recrystallized (ethyl acetate-diisopropyl ether) to give the titled compound (45 mg).

Melting point: 135 - 138°C

Reference Example 5

N-(4-Methoxyphenyl)-4-[[2-(2-piperidinoethyl)-6-tetralinyl]oxymethyl]benzamide



WSCD (0.11 ml) was added to DMF solution (2 ml) of 4-[[2-(2-piperidinoethyl)-6-tetralinyl]oxymethyl]benzoate (170 mg), 4-methoxyaniline (53 mg), HOBt (70 mg) and DMAP (60 mg) at room temperature, which was stirred for 12 hours. 10% aqueous potassium carbonate solution and water was added to the reaction mixture, and extraction was conducted using a mixed

WO 01/21577

PCT/JP00/06375


92

solution of THF and ethyl acetate. The organic layer was washed with water and saturated aqueous sodium chloride solution, dried, and then concentrated. The residue was purified using alumina column chromatography (development solvent: THF), and recrystallized (THF-IPE) to give the

5 titled compound (140 mg).
Melting point: 193 - 196°C

Reference Example 6

10 N-(3,4-Dimethoxyphenyl)-4-[[2-(2-piperidinoethyl)-6-tetralinyl]oxymethyl]benzamide




WSCD (free form, 0.2 ml) was added to DMF solution (3 ml) of 4-[[2-(2-piperidinoethyl)-6-tetralinyl]oxymethyl]benzoate (300 mg), 3,4-dimethoxyaniline (120 mg), HOBt (120 mg) and DMAP (100 mg) at room temperature, which was stirred for 12 hours. 10% aqueous potassium carbonate solution was added to the reaction mixture, and the resulting crystals were collected

20 by filtration. The crystals were washed with water, then dried. The crystals were purified using alumina column chromatography (development solvent; THF), and recrystallized (THF-IPE) to give the titled compound (330 mg).

25 Melting point: 178 - 180°C


Reference Example 7

6-[4-(Benzoylamino)benzyloxy]-2-(2-piperidinoethyl)tetralin



5 N-[4-(bromomethyl)phenyl]benzamide (670 mg) was added to the reaction mixture at room temperature, which was stirred for 1 hour. Water was added to the reaction mixture, and extraction was conducted using ethyl acetate. The organic layer was washed with water and saturated aqueous sodium chloride solution, dried, and then concentrated. The residue was purified using alumina column chromatography (development solvent : THF), and recrystallized (ethyl acetate) to give the titled compound (200 mg).
Melting point: 176 - 179°C

2-[(N,N-Dimethylamino)methyl]-6-tetralinyl 4-biphenylcarboxylate



20 4-Biphenylylcarboxylic acid (580 mg) and WSC (560 mg) were added to pyridine solution (6 ml) of 2-[(N,N-dimethylamino)methyl]-6-hydroxytetralin (300 mg), which was stirred at room temperature for 36 hours. Saturated sodium bicarbonate solution and water were added to the
25 reaction mixture, and extraction was conducted using ethyl acetate. The organic layer was washed with water and saturated aqueous sodium chloride solution, dried, and then

WO 01/21577

PCT/JP00/06375


94

concentrated. The residue was purified using alumina column chromatography (development solvent; hexane ~ hexane:ethyl acetate = 10:1), and recrystallized (hexane) to give the titled compound (300 mg).

5 Melting point: 85 - 86°C

Reference Example 9


2-[(N,N-Dimethylamino)methyl]-6-[4-[(4-methoxyphenyl)carbonyloxy]benzyloxy]tetralin



Diethyl azodicarboxylate (40% toluene solution, 950 mg) was added dropwise to THF solution (3 ml) of 2-[(N,N-dimethylamino)methyl]-6-hydroxytetralin (150 mg), 4-(hydroxymethyl)phenyl 4-methoxybenzoate (570 mg) and triphenylphosphine (574 mg) at room temperature, which was stirred for 3 hours. The reaction mixture was concentrated, and the residue was purified using alumina column chromatography (development solvent; hexane ~ hexane:ethyl acetate = 6:1), and recrystallized (ethyl acetate-hexane) to give the titled compound (175 mg).
Melting point: 119 - 121°C

Reference Example 10

25 2-[(N,N-Dimethylamino)methyl]-6-[4-[(4-methoxybenzyl)oxy]benzyloxy]tetralin



Diethyl azodicarboxylate (40% toluene solution, 1.91 g) was added dropwise to THF solution (6 ml) of 2-

WO 01/21577


PCT/JP00/06375

95

[(N,N-dimethylamino)methyl]-6-hydroxytetralin (300 mg),
4-[(4-methoxybenzyl)oxy]benzylalcohol (1.07 g) and
triphenylphosphine (1.15g) at room temperature, which was
stirred for 12 hours. The reaction mixture was
concentrated, and the residue was purified using alumina
column chromatography (development solvent; hexane -
hexane:ethyl acetate = 10:1), and recrystallized (ethyl
acetate-hexane) to give the titled compound (260 mg).
Melting point: 106 - 111°C

Reference Example 11

6-[4-[(1-Benzothiophen-2-yl)carbonylamino]benzyloxy]-2-
[(N,N-dimethylamino)methyl]tetralin



One drop of DMF was added to THF solution (4 ml) of
1-benzothiophene-2-carboxylic acid (230 mg), and oxalyl
chloride (0.23 ml) was further added under ice-cooling,
which was stirred for 30 minutes at room temperature. The
reaction mixture was concentrated, which was dissolved in
THF (1 ml). The obtained solution was added dropwise to
pyridine solution (6 ml) of 6-(4-aminobenzyloxy)-2-
[(N,N-dimethylamino)methyl]tetralin (300 mg), which was
stirred for 15 minutes. After stirring at room temperature
for another 15 minutes, 10% aqueous potassium carbonate
solution was added to the reaction mixture, and extraction
was conducted using ethyl acetate. The organic layer was
washed with water and saturated aqueous sodium chloride
solution, dried, and then concentrated. The residue was
purified using alumina column chromatography (development
solvent; ethyl acetate), and recrystallized (THF-IPE) to
give the titled compound (250 mg).
Melting point: 165 - 169°C


WO 01/21577

PCT/JP00/06375

96

Reference Example 12

2-[(N,N-Dimethylamino)methyl]-6-[4-[(4-methoxyphenyl)sulfonylamino]benzyloxy]tetralin




5

THF solution (1 ml) of 4-methoxybenzenesulfonyl chloride (270 mg) was added dropwise to pyridine solution (6 ml) of 6-[(4-aminobenzyl)oxy]-2-[(N,N-dimethylamino)methyl]tetralin (300 mg) under ice-cooling, which was stirred for 15 minutes. After stirring at room temperature for further 15 minutes, 10% aqueous potassium carbonate solution was added to the reaction mixture, and extraction was conducted using ethyl acetate. The organic layer was washed with water and saturated aqueous sodium chloride solution, dried, and then concentrated. The residue was purified using alumina column chromatography (development solvent: ethyl acetate), and recrystallized (ethyl acetate-IPE) to give the titled compound (260 mg). Melting point: 137 - 140°C

20

Reference Example 13

6-[4-(Benzylcarbonylamino)benzyloxy]-2-[(N,N-dimethylamino)methyl]tetralin



THF solution (1 ml) of phenylacetyl chloride (200 mg) was added dropwise to pyridine solution (6 ml) of 6-[(4-aminobenzyl)oxy]-2-[(N,N-dimethylamino)methyl]tetralin (300 mg) under ice-cooling, which was stirred for 15 minutes. After stirring at room temperature for further 15 minutes, saturated sodium

30

WO 01/21577


PCT/JP00/06375

97

bicarbonate solution was added to the reaction mixture, and extraction was conducted using ethyl acetate. The organic layer was washed with water and saturated aqueous sodium chloride solution, dried, and then concentrated. The residue was purified using alumina column chromatography (development solvent; hexane ~ hexane:ethyl acetate = 2:1), and recrystallized to give the titled compound (175 mg). Melting point: 130 - 135°C

Reference Example 14

6-[4-(Benzoylamino)benzyloxy]-2-[(N,N-dimethylamino)methyl] tetralin



Benzoyl chloride (0.14 ml) was added dropwise to pyridine solution (6 ml) of 6-[(4-aminobenzyl)oxy]-2-[(N,N-dimethylamino)methyl]tetralin (300 mg) under ice-cooling, which was stirred at room temperature for 30 minutes. 10% aqueous potassium carbonate solution was added to the reaction mixture, and extraction was conducted using ethyl acetate. The organic layer was washed with water and saturated aqueous sodium chloride solution, dried, and then concentrated. The residue was purified using alumina column chromatography (development solvent; ethyl acetate), and recrystallized (THF-IPE) to give the titled compound (240 mg). Melting point: 128 - 133°C


Reference Example 15

2-[(N,N-Dimethylamino)methyl]-6-[4-[(4-methoxybenzoyl)amino]benzyloxy]tetralin

WO 01/21577

PCT/JP00/06375

98




p-Anisoyl chloride (0.20 ml) was added dropwise to pyridine solution (6 ml) of 6-[(4-aminobenzyl)oxy]-2-[(N,N-dimethylamino)methyl]tetralin (300 mg) under ice-cooling, which was stirred at room temperature for 30 minutes. 10% aqueous potassium carbonate solution was added to the reaction mixture, and extraction was conducted using ethyl acetate. The organic layer was washed with water and saturated aqueous sodium chloride solution, dried, and then concentrated. The residue was purified using alumina column chromatography (development solvent: ethyl acetate), and recrystallized (THF-IPE) to give the titled compound (300 mg).
Melting point: 155 - 159°C

15

Reference Example 16

2-[(N,N-Dimethylamino)methyl]-6-[4-[(2-methoxybenzoyl)amino]benzyloxy]tetralin



o-Anisoyl chloride (0.15 ml) was added dropwise to pyridine solution (4 ml) of 6-[(4-aminobenzyl)oxy]-2-[(N,N-dimethylamino)methyl]tetralin (200 mg) under ice-cooling, which was stirred at room temperature for 30 minutes. 10% aqueous potassium carbonate solution was added to the reaction mixture, and extraction was conducted using ethyl acetate. The organic layer was washed with water and saturated aqueous sodium chloride solution, dried, and then concentrated. The residue was purified

WO 01/21577

PCT/JP00/06375

99


using alumina column chromatography (development solvent; THF), and recrystallized (ethyl acetate-hexane) to give the titled compound (200 mg).

Melting point: 106 - 108°C

5

Reference Example 17

6-[4-[N-(4-Methoxybenzoyl)-N-methylamino]benzyloxy]-2-[(N,N-dimethylamino)methyl]tetralin



- 10 Diethyl azodicarboxylate (40% toluene solution, 960 mg) was added dropwise to THF solution (3 ml) of 2-[(N,N-dimethylamino)methyl]-6-hydroxytetralin (150 mg), N-[4-(hydroxymethylphenyl)-4-methoxy-N-methylbenzamide (600 mg) and triphenylphosphine (570 mg) at room
- 15 temperature, which was stirred for 12 hours. After the reaction mixture was concentrated, the residue was purified using silica gel column chromatography (development solvent; hexane ~ ethyl acetate ~ ethyl acetate:methanol = 1:2), and then purified using alumina column
- 20 chromatography (development solvent; hexane ~ hexane:ethyl acetate = 2:1) to give the titled compound (185 mg).
¹H-NMR (CDCl₃) δ: 1.20-1.50(1H, m), 1.80-2.46(5H, m), 2.25(6H, s), 2.68-2.86(3H, m), 3.47(3H, s), 3.74(3H, s), 4.95(2H, s), 6.52-6.76(4H, m), 6.84-7.14(3H, m), 7.22-
 25 7.38(4H, m).

Reference Example 18

N-[4-[[[2-(Diethylamino)ethyl]amino]carbonyl]phenyl] 4-biphenylylcarboxamide